

A novel method to induce solvent tolerance in bacteria

Jamie Hinks^{1*}, Yaofeng Wang², Artur Matysik², Muguang Yu², Stefan Wuertz^{1,3}, Thomas Seviour^{1*}

Biobutanol has vast potential as a drop-in alternative to fossil fuel (1, 2). Butanol can be produced biologically by Clostridium beijerinckii and Clostridium acetobutylicum although biological production is uneconomical due to low process yields which account for high recovery costs by distillation (2-4). Poor microbiological solvent tolerance underlies poor process yields in biological solventogenesis, including butanol production. Solvents are known to compromise the integrity of biological membrane, accordingly research endeavours to develop solvent tolerant microbes have focused on shoring up membrane integrity in instances of solvent challenge using engineering, metabolic and adaptive approaches (2, 5-10).

Membrane Insertion Molecules (MIMs) are molecules that modify cell function or stability by intercalating into the phospholipid bilayer of microbial membranes. There are two types of MIM; those designed to act as antimicrobial agents, killing cells by destabilizing the lipid bilayer, and enhancement MIMs, which are designed to improve cell function (11).

We show that a proprietary MIM can increase butanol tolerance in E. coli K12 which, when treated, can outgrow an untreated control in concentrations up to 3.5% v/v butanol. Using a combination of molecular dynamic modelling and single molecule tracking in model membranes treated with MIM_{ST}, along with membrane leakage and cell/butanol partitioning assays, we propose a mechanism for how MIM_{ST} treatment may increase solvent tolerance in microbes. We discuss the benefits and shortcomings of this technique and the relationship between solvent tolerance and solvent yield in solventogenesis.

Keywords—butanol, tolerance, biofuels, membranes, modification

Jamie Hinks Thomas Seviour, Stefan Wuertz

1 Singapore Centre on Environmental Life Science Engineering(SCELSE),
Nanyang Technological University,

Yaofeng Wang, Artur Matysik, Yuguang Mu

2 School of Biological Sciences (SBS), Nanyang Technological University,
Singapore

Stefan Wuertz

3 Department of Civil and Environmental Engineering, University of
California Davis, Davis, CA 95616, USA

Acknowledgments

SCELSE is funded by Singapore's National Research Foundation, Ministry of Education, Nanyang Technological University (NTU) and National University of Singapore (NUS), and hosted by NTU in partnership with NUS.

References

- [1] 1. Savage N. 2011. Fuel options: The ideal biofuel. Nature 474:S9-S11.
- [2] 2. Qureshi N, Blaschek H. 2000. Economics of Butanol Fermentation using Hyper-Butanol Producing Clostridium Beijerinckii BA101. Food and bioproducts processing 78:139-144.
- [3] 3. Baer SH, Blaschek HP, Smith TL. 1987. Effect of butanol challenge and temperature on lipid composition and membrane fluidity of butanol-tolerant Clostridium acetobutylicum. Appl. Environ. Microbiol. 53:2854-2861.
- [4] 4. Zheng Y-N, Li L-Z, Xian M, Ma Y-J, Yang J-M, Xu X, He D-Z. 2009. Problems with the microbial production of butanol. J. Ind. Microbiol. Biotechnol. 36:1127-1138.
- [5] 5. Kanno M, Katayama T, Tamaki H, Mitani Y, Meng X-Y, Hori T, Narihiro T, Morita N, Hoshino T, Yumoto I, Kimura N, Hanada S, Kamagata Y. 2013. Isolation of Butanol- and Isobutanol-Tolerant Bacteria and Physiological Characterization of Their Butanol Tolerance. Appl. Environ. Microbiol. 79:6998-7005.
- [6] 6. Zhang H, Chong H, Ching C, Song H, Jiang R. 2012. Engineering global transcription factor cyclic AMP receptor protein of Escherichia coli for improved 1-butanol tolerance. Appl. Microbiol. Biotechnol. 94:1107-1117.
- [7] 7. Huffer S, Clark ME, Ning JC, Blanch HW, Clark DS. 2011. Role of Alcohols in Growth, Lipid Composition, and Membrane Fluidity of Yeasts, Bacteria, and Archaea. Appl. Environ. Microbiol. 77:6400-6408.
- [8] 8. Knoshaug E, Zhang M. 2009. Butanol Tolerance in a Selection of Microorganisms. Appl. Biochem. Biotechnol. 153:13-20.
- [9] 9. Isken S, Derks A, Wolffs PF, de Bont JA. 1999. Effect of organic solvents on the yield of solvent-tolerant Pseudomonas putida S12. Appl. Environ. Microbiol. 65:2631-2635.
- [10] 10. Rühl J, Schmid A, Blank LM. 2009. Selected Pseudomonas putida Strains Able To Grow in the Presence of High Butanol Concentrations. Appl. Environ. Microbiol. 75:4653-4656.
- [11] Hill EH, Stratton K, Whitten DG, Evans DG. 2012. Molecular Dynamics Simulation Study of the Interaction of Cationic Biocides with Lipid Bilayers: Aggregation Effects and Bilayer Damage. Langmuir 28:14849-14854.