

# The existence of positive solutions for three-point boundary value problems of fractional differential equations

Li Zou, Shenjie Fan, Jiushuan Zhang, Lizheng Xue, Dong Pan, Shurong Sun\*

**Abstract**—In this paper, we investigate the existence of solutions for a class of three-point boundary value problems of fractional differential equations. By using Lebesgue dominated convergence theorem and Schauder fixed point theorem, a sufficient condition of the existence solution is obtained.

**Keywords**—fixed point theorem; dominated convergence theorem; the existence of solutions

## 1. Introduction

Fractional calculus, the study of integration and differentiation of fractional order, has recently been extended to include its discrete analogues of fractional difference calculus and fractional quantum calculus. Due to the similarities of the three theories there has been research on whether there exist a single theory that encapsulates them.

Fractional differential equations are extension and expansion of integer-order differential equations. At present, the fractional differential equations have been increasingly applied to different fields of sciences.

Fractional differential equations are using widely in the fields of various sciences such as physics, mechanic, chemistry, engineering. Fractional differential equations have been of increasing importance for the past decades due to their diverse applications in science and engineering, such as, the memory of a variety of materials, signal identification and image processing, optical systems, thermal system materials and mechanical systems, control system. Many interesting results of the existence of solutions of various classes of fractional differential equations have been obtained, see [1-24] and the references therein.

---

Li Zou, Shenjie Fan, Jiushuan Zhang, Lizheng Xue, Dong Pan, School of Mathematical Sciences, University of Jinan, Jinan, Shandong 250022, P R China.

\*Corresponding author: Shurong Sun, School of Mathematical Sciences, University of Jinan, Jinan, Shandong 250022, P R China.

Many scholars paid more attention to it, and there many interesting results on the existence of solutions of various classes of fractional differential equations have been obtained.

Recently, much attention has been focused on the study of the existence and multiplicity of solutions or positive solutions for boundary value problems of fractional differential equations with boundary value problems by the use of techniques of nonlinear analysis (fixed-point theorems, Leray-Schauder theory, the upper and lower solution method).

Boundary value problems of the fractional differential equations are branches of the fractional differential equations and have received much attention. There have been many results for boundary value problems of the fractional differential equations, such as the solutions for boundary value problem of fractional differential equations and m-point boundary value problems [1].

Gan in [18] investigated the existence and non-existence of positive solution for contains the parameters of two-point boundary value problems of fractional differential equations by using fixed point theorem on compression and expansion of cones. Liang and Zhang in [19] deals with the existence of positive solutions for a class of two-point boundary value problems of fractional differential equations by means of upper and lower solutions method and fixed point theorem. Xu, Xian and Zhou in [20] investigate the existence of solutions for a class of periodic boundary value problems of fractional differential equations by using contraction mapping principle.

Inspired by the above document, we investigate the following the existence of solutions of fractional differential equations.

$$\begin{cases} D_{0+}^{\alpha} u(t) = f(t, u(t)), & t \in J = [0, T], \\ u(0) = 0, u(T) = b u(\eta), \end{cases} \quad (1)$$

where  $1 < \alpha \leq 2$ ,  $0 < b < 1$ ,  $\eta \in (0, T)$ ,  $f : J \times \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ , this paper provides a sufficient condition of the existence of solution for the problem (1) by using Schauder fixed point theorem.



## 2. Preliminaries

**Definition 2.1.** The fractional integral of order  $\alpha$  for a function  $y : (0, \infty) \rightarrow R$  is defined as:

$$I_{0+}^{\alpha} y(t) = \frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} y(s) ds,$$

where  $\alpha > 0$ ,  $\Gamma(\cdot)$  denotes the Gamma function.

**Definition 2.2.** [1] For a continuous function  $y : (0, \infty) \rightarrow R$ ,

the Caputo derivative of fractional order  $\alpha$  is defined as

$$D_{0+}^{\alpha} y(t) = \frac{1}{\Gamma(n-\alpha)} \left(\frac{d}{dt}\right)^n \int_0^t (t-s)^{n-\alpha-1} y^{(n)}(s) ds,$$

where  $\alpha > 0$ ,  $n = [\alpha] + 1$ ,  $[\alpha]$  denotes the integer part of the real number  $\alpha$ ,  $\Gamma(\cdot)$  denotes the Gamma function.

**Lemma 2.1.** [1] Let  $\alpha > 0$ , if  $u \in C(0,1) \cap L^1(0,1)$ ,  $D_{0+}^{\alpha} u(t) \in C(0,1) \cap L^1(0,1)$ , for some  $c_i \in \mathfrak{R}, i=1,2,3,\dots,N$ , subject to

$$I_{0+}^{\alpha} {}^c D_{0+}^{\alpha} u(t) = u(t) + c_1 + c_2 t + \dots + c_N t^{n-1},$$

$$c_i \in \mathfrak{R}, i=1,2,\dots,n,$$

where  $n = [\alpha] + 1$ .

**Lemma 2.2.** Let  $h \in C(J, R)$ , a function  $u$  is a solution of the fractional BVP

$$\begin{cases} {}^c D_{0+}^{\alpha} u(t) = h(t), & t \in J, 1 < \alpha \leq 2, \\ u(0) = 0, u(T) = b u(\eta). \end{cases} \quad (2)$$

If and only if  $u$  is a solution of the fractional integral equation

$$\begin{aligned} u(t) = & \frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} h(s) ds \\ & + \frac{ct}{\Gamma(\alpha)} \int_0^T (T-s)^{\alpha-1} h(s) ds \\ & - \frac{cbt}{\Gamma(\alpha)} \int_0^{\eta} (\eta-s)^{\alpha-1} h(s) ds, \end{aligned}$$

where  $c = \frac{1}{\eta - T}$ .

**Proof:** By Lemma 2.1, transform the problem (2) into an integral equation

$$u(t) = I_{0+}^{\alpha} h(t) + c_1 + c_2 t, \quad c_1, c_2 \in R,$$

by the boundary value conditions

$$u(0) = 0, u(T) = b u(\eta),$$

we can get

$$\begin{cases} c_1 = 0, \\ c_2 = \frac{ct}{\Gamma(\alpha)} \int_0^T (T-s)^{\alpha-1} h(s) ds - \frac{cbt}{\Gamma(\alpha)} \int_0^{\eta} (\eta-s)^{\alpha-1} h(s) ds. \end{cases}$$

where  $c = \frac{1}{\eta - T}$ . Thus

$$\begin{aligned} u(t) = & \frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} h(s) ds \\ & + \frac{ct}{\Gamma(\alpha)} \int_0^T (T-s)^{\alpha-1} h(s) ds \\ & - \frac{cbt}{\Gamma(\alpha)} \int_0^{\eta} (\eta-s)^{\alpha-1} h(s) ds. \end{aligned}$$

They are equivalent.

Measurable function  $m : J \rightarrow R$ , defined norm

$$\|m\|_{L^p(J, R)} = \begin{cases} \left( \int |m(t)|^p dt \right)^{\frac{1}{p}}, & 0 \leq p < \infty, \\ \inf_{\mu(J)} \left\{ \sup_{t \in J} |m(t)| \right\}, & p = \infty. \end{cases}$$

where  $\mu(J)$  is Lebesgue measure on  $J$ ,  $L^p(J, R)$  is a Banach space consists of all Lebesgue measurable function  $m : J \rightarrow R$ ,  $\|m\|_{L^p(J, R)} < \infty$ .

**Lemma 2.3.** [1] Lebesgue dominated convergence theorem:

Supposes that  $E \subseteq R^q$  is measurable set,  $\{f_n\}_{n=1}^{\infty}$  is a sequence of measurable function on  $E$ .  $F$  is nonnegative  $L$  integrable function on  $E$ , if for any natural number  $n$ ,

$$\begin{aligned} |f_n(x)| & \leq F(x) \text{ a.e on } E, \\ \lim_{n \rightarrow \infty} f_n(x) & = f(x), \end{aligned}$$

thus

$$\begin{aligned} \lim_{n \rightarrow \infty} \int_E |f_n(x) - f(x)| dx & = 0, \\ \lim_{n \rightarrow \infty} \int_E f_n(x) dx & = \int_E f(x) dx. \end{aligned}$$

**Lemma 2.4.** [1] Schaefer fixed point theorem. Supposes  $X$  is a Banach space, mapping  $T : X \rightarrow X$  is a completely continuous mapping. If set

$$V = \left\{ u \in X \mid u = \mu T u, 0 \leq \mu < 1 \right\}$$

is a bounded set, then mapping  $T$  has a fixed point in  $X$ .

## 3. Main Results

Definition:  $X = C(J, R)$ , its norm is  $\|u\| = \sup_{t \in J} |u(t)|$ , it is obvious that  $X$  is a Banach space. We provide a sufficient condition of the existence of solution

for the problem (1) by Schaefer fixed point theorem. We make the following conditions through out this paper:

$$(H_1) f : J \times R \times R \rightarrow R \text{ is continuous}$$

(H<sub>2</sub>) there exists a constant  $L_1 > 0$ , for  $\forall t \in J$  and for all  $u \in X$ ,

$$|f(t, u(t))| \leq L_1.$$

**Theorem 3.1** Assuming the conditions of the (H<sub>1</sub>) and (H<sub>2</sub>) are satisfied. Then the BVP (1) has a positive solutions at least.

**Proof:** By Lemma 2.2, the promble (1) is equivalent to

$$u(t) = \frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} f(s, u(s)) ds + \frac{ct}{\Gamma(\alpha)} \int_0^T (T-s)^{\alpha-1} f(s, u(s)) ds - \frac{cbt}{\Gamma(\alpha)} \int_0^\eta (\eta-s)^{\alpha-1} f(s, u(s)) ds.$$

Considering the mapping  $F : X \rightarrow X$  is followed.

$$(Fu)(t) = \frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} f(s, u(s)) ds + \frac{ct}{\Gamma(\alpha)} \int_0^T (T-s)^{\alpha-1} f(s, u(s)) ds - \frac{cbt}{\Gamma(\alpha)} \int_0^\eta (\eta-s)^{\alpha-1} f(s, u(s)) ds,$$

therefore, the existence of positive solutions for three-point boundary value problems (1) are turning to problems of the existence of fixed point that mapping  $F$  in  $X$ .

Fristly, proving  $F$  is continuous.

Take the series  $\{u_n\} \subset X$ , let the  $u_n \rightarrow u (n \rightarrow \infty)$ , by the Lebesgue dominated convergence theorem, we can get  $\forall t \in J$ , there is

$$\begin{aligned} & |(Fu_n)(t) - (Fu)(t)| \\ & \leq \frac{1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} \|f(s, u_n(s)) - f(s, u(s))\| ds \\ & + \frac{ct}{\Gamma(\alpha)} \int_0^T (T-s)^{\alpha-1} \|f(s, u_n(s)) - f(s, u(s))\| ds \\ & + \frac{cbt}{\Gamma(\alpha)} \int_0^\eta (\eta-s)^{\alpha-1} \|f(s, u_n(s)) - f(s, u(s))\| ds \\ & \leq \frac{T^\alpha + (1+b)cT^{\alpha+1}}{\Gamma(\alpha+1)} \|f(s, u_n(s)) - f(s, u(s))\|, \end{aligned}$$

by the conditions of the (H<sub>1</sub>) :  $f$  is continuous, so

$$\|f(s, u_n(s)) - f(s, u(s))\| \rightarrow 0,$$

therefore we can draw a conclusion that  $F$  is continuous.

And then, we prove that  $F$  mapped a bounded set to a bounded set in  $X$ .

Let  $\Omega \subset X$  is a bounded set.  $\forall u \in \Omega$ , Combining with (H<sub>2</sub>), we have

$$\begin{aligned} |(Fu)(t)| & \leq \frac{L_1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} ds \\ & + \frac{cL_1t}{\Gamma(\alpha)} \int_0^T (T-s)^{\alpha-1} ds + \frac{cL_1bt}{\Gamma(\alpha)} \int_0^\eta (\eta-s)^{\alpha-1} ds \\ & \leq \frac{L_1t^\alpha}{\alpha\Gamma(\alpha)} + \frac{cL_1T^\alpha}{\alpha\Gamma(\alpha)} + \frac{cbL_1\eta^\alpha}{\alpha\Gamma(\alpha)} \\ & \leq \frac{L_1T^\alpha + (1+b)L_1cT^{\alpha+1}}{\Gamma(\alpha+1)} = L_2. \end{aligned}$$

Namely, there is constant  $L_2 > 0$ , make  $\|Fu\| \leq L_2$ , thus, we know that  $F$  mapped a bounded set to a bounded.

Finally, we will prove that the mapping  $F$  is equicontinuous.

Set  $0 < t_1 < t_2 < T$ , for  $\forall u \in X$ , by (H<sub>2</sub>) we have

$$\begin{aligned} & |(Fu)(t_2) - (Fu)(t_1)| \\ & \leq \frac{1}{\Gamma(\alpha)} \left| \int_0^{t_2} (t-s)^{\alpha-1} f(s, u(s)) ds - \int_0^{t_1} (t-s)^{\alpha-1} f(s, u(s)) ds \right| \\ & + \frac{c(t_2-t_1)}{\Gamma(\alpha)} \int_0^T (T-s)^{\alpha-1} |f(s, u(s))| ds \\ & + \frac{cb(t_2-t_1)}{\Gamma(\alpha)} \int_0^\eta (\eta-s)^{\alpha-1} |f(s, u(s))| ds \\ & \leq \frac{L_1}{\Gamma(\alpha)} (t_2^\alpha - t_1^\alpha) + \frac{c\eta^\alpha L_1 (t_2 - t_1)}{\alpha\Gamma(\alpha)} + \frac{cT^\alpha L_1 b (t_2 - t_1)}{\Gamma(\alpha)} \\ & \leq \frac{L_1}{\Gamma(\alpha+1)} (t_2^\alpha - t_1^\alpha) + \frac{c(1+b)T^\alpha L_1 (t_2 - t_1)}{\Gamma(\alpha+1)}, \end{aligned}$$

when  $t_1 \rightarrow t_2$ , we have

$$|(Fu)(t_2) - (Fu)(t_1)| \rightarrow 0,$$

so  $F$  is equicontinuous on  $J$ . According to Ascoli-Arzela theorem,  $F$  is completely continuous.

Meanwhile, we prove that set

$$V = \{u \in X, u = \mu Fu, 0 < \mu < 1\}$$

is bounded set.

Let  $u \in V$ , then,  $u = \mu Fu, 0 < \mu < 1$ , for  $\forall t \in J$ , we have

$$\begin{aligned} |u(t)| & = |\mu(Fu)(t)| \leq \frac{L_1}{\Gamma(\alpha)} \int_0^t (t-s)^{\alpha-1} ds \\ & + \frac{cL_1t}{\Gamma(\alpha)} \int_0^T (T-s)^{\alpha-1} ds + \frac{cL_1bt}{\Gamma(\alpha)} \int_0^\eta (\eta-s)^{\alpha-1} ds \\ & \leq \frac{L_1T^\alpha + 2cL_1T^{\alpha+1}}{\Gamma(\alpha+1)} = L_3. \end{aligned}$$

There is a constant  $L_3 > 0$ , make  $\|u\| \leq L_3$ , then we know that the set  $V$  is bounded set.

Therefore, in summary, schauder fixed point theorem shows,  $F$  in  $X$  has at least one fixed point, the boundary conditions (1) has at least one solution.

This proves that the existence of solutions with the boundary condition (1).

### Acknowledgment

This research is supported by Shandong Provincial Natural Science Foundation (ZR2012AM009, ZR2013AL003).

### References

- [1] Y I Podlubn., Fractional Differential Equation, Mathematics in Science and Engineering. New York, London, Toronto: Academic Press,1999.
- [2] S. Sun, Q. Li, Y. Li, Existence and uniqueness of solutions for a coupled system of multi-term nonlinear fractional differential equations, *Comput Math Appl*, 64 (2012) 3310–3320.
- [3] Y. Zhao, S. Sun, Z. Han, M.Zhang, Positive solutions for boundary value problems of nonlinear fractional differential equations, *Appl Math Comput*, 217 (2011) 6950–6958.
- [4] Y. Zhao, S. Sun, Z. Han, Q. li, Positive Solutions to Boundary Value Problems of Nonlinear Fractional Differential Equations, *Abs Appl Anal*, 2011 (2011), Article ID 390543, 1–16.
- [5] Y. Zhao, S. Sun, Z. Han, Q. Li, The existence of multiple positive solutions for boundary value problems of nonlinear fractional differential equations, *Commun Nonlinear Sci Numer Simulat*, 16 (2011) 2086–2097.
- [6] W. Feng, S. Sun, Z. Han, Y. Zhao, Existence of solutions for a singular system of nonlinear fractional differential equations, *Comput Math Appl*, 62 (2011) 1370–1378.
- [7] S. Sun, Y. Zhao, Z. Han, Y. Li, The existence of solutions for boundary value problem of fractional hybrid differential equations, *Commun Nonlinear Sci Numer Simulat*,17 (2012) 4961–4967.
- [8] S. Sun, Y. Zhao, Z. Han, M. Xu, Uniqueness of positive solutions for boundary value problems of singular fractional differential equations, *Inverse Probl Sci Eng*,20 (2012) 299–309.
- [9] S. Sun, Y. Zhao, Z. Han, J.Liu. Eigenvalue problem for a class of nonlinear fractional differential equations, *Annals of Functional Analysis*, 4(1) (2013) 25–39.
- [10] Y. Li, S. Sun, D. Yang, Z. Han, Three point boundary value problems of fractional functional differential equations with delay, *Boundary Value Problems*, 2013(38) (2013) 1–15.
- [11] Y. Li, S. Sun, Z. Han, H. Lu, The existence of positive solutions for boundary value problem of fractional Sturm-Liouville functional differential equation. *Abstract and Applied Analysis*,2013, (2013) Article ID 301560, 1–20.
- [12] X. Li, Z. Han, S. Sun, Existence of positive solutions of nonlinear fractional q-difference equation with parameter. *Advances in Difference Equations*,2013(260) (2013)1–13.
- [13] Y. Pan, Z. Han, S. Sun, C. Hou, The existence of solutions to a class of boundary value problems with fractional difference equations, *Advances in Difference Equations*, 2013 (275) (2013) 1–20.
- [14] R. Yan, S. Sun, H. Lu, Y. Zhao, Existence of solutions for fractional differential equations with integral boundary conditions, *Advances in Difference Equations*, 2014(25) (2014) 1–13.
- [15] H. Lu, Z. Han, S. Sun, Multiplicity of positive solutions for Sturm-Liouville boundary value problems of fractional differential equations with p-Laplacian. *Boundary Value Problems*, 2014 (26) (2014) 1–17.
- [16] Y. Wang, Z. Han, P. Zhao, S. Sun, On the oscillation and asymptotic behavior for a kind of fractional differential equations, *Advances in Difference Equations*, 2014 (50) (2014) 1–11.
- [17] X. Li, Z. Han, S. Sun, H. Lu, Boundary value problems for fractional  $q$ -difference equations with nonlocal conditions, *Advances in Difference Equations*, 2014(57) (2014) 1–16.
- [18] S. Q.Gan, Dissipativity of methods for nonlinear Volterra delay integro differential equations, *J Comput Math*, 206 (2007)98–107.
- [19] S.Liang, J.Zhang, Positive solutions for boundary value problems of nonlinear fractional differential equation., *Nonlinear Anal*, 71(2009)5545–5550.
- [20] J. Xu, Z. Xiao, W. Zhou, The existence of solutions for 2K order ordinary differential equation, *Journal of Yunnan University: Natural Science Edition*, 33 (4) (2011) 373–377.
- [21] C.S. Goodrich, Existence and uniqueness of solutions to a fractional difference equation with non-local conditions, *Comput Math Appl*, 61(2011) 191–202.
- [22] Z. Han, Y. Zhao, Y. Sun, C. Zhang, Oscillation for a class of fractional differential equation, *Discrete Dynamics in Nature and Society*,2013(2013) Article ID 390282, 1–6.
- [23] S. Sun, Q. Li, Y. Li, Existence and uniqueness of solutions for a coupled system of multi-term nonlinear fractional differential equations, *Comput Math Appl*, 64 (2012) 3310–3320.
- [23] Q. Li, S. Sun, P. Zhao, Z. Han, Existence and uniqueness of solutions for initial value problem of nonlinear fractional differential equations, *Abstract and Applied Analysis*, 2012 (2012) Article ID 615230, 1–14.
- [24] Y. Pan, Z. Han, S. Sun, Y. Zhao, The existence of solutions to a system of discrete fractional boundary value problems, *Abstract and Applied Analysis*, 2012 (2012) Article ID 707631, 1–15.