

Assessment of Microbial Community and Soil Enzyme Activity of Coal Mine Dumps of Sonbhadra Uttar Pradesh, India.

Shobhit Verma, Abhay Pratap Singh, Shikha Devi, Radha Rani Mewaram, Shivesh Sharma*, Kumud Dubey

Abstract— Mining activities alter the normal soil properties and negatively affect the ecosystem and environment by many ways. Assessing the damage done to the soil is very important to know the actual status of soil health. Soil health and fertility can be measured by the amount of microorganisms present and activity of different soil enzymes. Number of microorganisms shows amount of stress and unfavorable conditions. In the present study, colony forming units of microorganisms were found relatively higher in natural forest soils in comparison to soil collected from dump sites. Soil enzyme activities are very sensitive to both natural and anthropogenic disturbances and show a quick response to the induced changes so they can express the changes that may have happened in the soil due to mining. Soil phosphatase and dehydrogenase activity in natural forest have been found higher as compared to dump soil. In contrast, higher catalase activity was observed in dump soil and lower in Natural Forest soil.

Keywords—Mining, Soil Health, microorganisms, soil enzyme activities.

I. Introduction

Soil provides a medium where a long-term interaction among microorganisms, soil minerals and organic matter can influence the physio-chemical and biological characteristics of terrestrial system. Mining activity, specifically open cast mining, often lead to land degradation with adverse changes in soil textural and structural attributes (1). Open cast coal mining leads to the removal of overlying soil and formation of heaps of rocks, coal seam, sand, minerals and soil with very low nutrient quantity (2, 3). The Open cast mining thus leads to a nutrient deficient condition for soil microorganisms and plants, and poses problem for the process of revegetation (2, 4), restoration (5, 6) and pedogenesis (7, 8). Heavy metals are dangerous group of pollutants, which cannot be naturally degraded like organic pollutants and they accumulate in different parts of the food chain (9). Soil Enzymes are produced by microorganisms, either as a extracellular

secretion or intracellular product released by lysis of cell, which provides an insight into microbial activity and dynamics (10). Biogeochemical cycles are regulated by the microbial activity in the soil or by the enzymes they produce (11). These enzymes are constantly synthesized, accumulated, inactivate/decomposed in the soil, and during this they stabilize the soil structure and function by nutrient mineralization and cycling, transformation and energy metabolism, decomposition and formation of organic matter (12), and decomposition of xenobiotics. Dumps of mining are also a threat to the surrounding environment as it contains several trace elements in high amounts (13, 14). The problem of coal mine dumps and overburden generated by opencast mining is enormous in India. Presently, the north eastern coalfields of Coal India Limited (NECF-CIL), Margherita, Assam, have produced more than 1000 ha of mine OB wasteland. Reports are also available on the prospects and environmental issues related to the north east (NE) collieries (15, 16).

The present study aims to find out the effect of mining over the environment and ecosystem utilizing microorganisms and soil enzyme activity as possible indicators.

II. Materials and Method

A. Study site and Soil Sampling

Sonbhadra is the second largest district of Uttar Pradesh India, with the area of 6788 km². Sonbhadra is located in the south-eastern ranges of Vindhyanal mountains which are having treasures of minerals, metals and most importantly Coal. There are various coal mines in Sonbhadra out of which Kakari mine was selected for soil sample collection. The Kakari mine is located (Lat 24°16'55"- 24°18'90"N and Long 82°74'25"- 82°76'72"E) beside Govind Ballabh Pant Sagar. Kakri mines are developed under kakri project of Northern Coal fields limited a subsidiary of Coal India Limited.

Soil samples were collected from various dumps of the mine as well as from the nearby forest. Three samples in triplicate were collected from different spots at dump site (D1, D2 & D3) and same number of samples was taken from the nearby natural forest (NF1, NF2& NF3).

B. Soil pH estimation

For estimation of soil pH solution of soil and water was prepared in the ratio of 1:2 and incubated for half an hour

Shobhit Verma, Abhay Pratap Singh, Shikha Devi, Radha Rani Mewaram, Shivesh Sharma*

Department of Biotechnology, Motilal Nehru National Institute of Technology Allahabad, India

Kumud Dubey

Centre for Social Forestry and Eco-rehabilitation Allahabad, India

at 30°C and the pH was measured with the help of electrode of pH meter.

C. Microbial Enumeration

For counting the colony forming units, samples were serially diluted with 0.85% saline solution and plated on Nutrient agar (Hi-media) plates. After incubation of about 30hrs. colony forming units of different plates were counted with the help of colony counter, calculated and expressed in CFU/g.(17).

D. Soil Enzyme Activities

As soil enzyme activities are the indirect measures of the microbial activity in the soil, different soil enzymes named catalase, dehydrogenase and phosphatase were tested.

Catalase enzyme activity was assessed by titrimetric method using KMnO_4 (18). 2 g of soil mixed with 40 ml of water and 5 ml of 0.3% H_2O_2 added to it and shaken for 20 minutes at 25°C. Now solution was filtered and added 5ml of 1.5 mol/l H_2SO_4 to 10 ml of the filtrate, remaining H_2O_2 was measured by titrating it by 0.1 mol/l KMnO_4 solution. The activity was expressed as ml 0.1 mol/l KMnO_4 sol. titrated/(g dry soil x 20 min.)

Dehydrogenase activity was measured by the reduction of 2,3,5-triphenyltetrazolium chloride (TTC) (19, 20). For this 1.0g of soil was weighed and kept in a clean test tube and added 1.0 ml of TTC then incubated for 96 hours at 27°C, after which 10 ml of methanol was added to extract the triphenyl formazon(TPF) formed during the reaction. Amount of TPF released was determined by spectrophotometric analysis at 485nm.

For measuring phosphatase activity, 1.0g of soil sample was taken in a test tube and 5ml of 100.0mM phosphate buffer was included. 10.0mM of p-nitro phenyl phosphate (p-NPP) in 1.0ml solution was used as substrate. The final volume of the reaction mixture was adjusted to 10.0ml with the addition of requisite amount of distilled water. The tube was vortexed (2.0min.) at room temperature. It was then incubated at 37°C for 60min in shaker (100 rpm). After the incubation period, the sample was centrifuged at 10000 rpm for 5.0min. The clear supernatant was taken in a clean test tube and 10.0ml of 1.0M NaOH was added. The yellow colored filtrate was analyzed using a colorimeter at $\lambda=430\text{nm}$ (21).

III. Results and Discussion

As the soil samples were subjected for the pH testing and it was found that the Dump 1(D1) is most acidic (pH 5.26) among all and NF 3 is most basic (pH 6.82). All the dump soil samples showed lower pH values which indicated towards higher acidity while all natural forest soils showed higher pH value than dump soils (Graph 1). The pH value of soil is often modeled as a positive liner relationship with soil fertility and productivity, where a high pH indicates a better soil. Improving soil chemical condition by the reduction of soil


acidity has been well studied (1) Different studies have proved that due to excessive mineral deposition during mining soil turns acidic (2, 22). Improvement of soil pH due to natural succession or by plantation strategy on coal mine spoil has been reported (2, 23).

The colony forming units when calculated revealed that all natural forest soil samples(NF1,NF2 & NF3) have much more colony forming units of microbes than the dump soil of Kakari mines, which proved that number of microorganisms substantially decrease in the dump soil.(Graph 2)


Soil catalase activity was found to be higher in the dump soil than in natural forest soils, and the highest catalase activity was measured in D2 that is 0.0075 ml of .1 M KMnO_4 g^{-1} of soil min^{-1} , while minimum 0.00375 ml of .1 M KMnO_4 g^{-1} of soil min^{-1} showed by NF1. The high activity of catalase in dump soils shows high stress condition for microorganisms (18).

Results of soil dehydrogenase activity assay showed the highest dehydrogenase activity in natural forest soil sample (NF 2) i.e. 1.02 $\mu\text{g.g}^{-1}\text{dry soil.h}^{-1}$ and lowest activity in dump soil sample (D2) 0.1625 $\mu\text{g.g}^{-1}\text{dry soil.h}^{-1}$. Over all natural forest soil showed much higher dehydrogenase activity than dump soils. Estimation of dehydrogenase activity is very essential due to the fact that they are an integral part of soil microorganisms and are involved electron transport system, and requires viable cells to express its activity (24). Dehydrogenase is considered to be an index of microbial activity and metabolic status of soil microorganisms (25, 26, 27, 28). The results proved that microbial activity is higher in natural forest soil.


Phosphatase activity appears to be more affected by the metabolic state of soil, biological activities of microbial population, so their activity level can be used as an index for microbial activity in soil (29). Highest amount of phosphatase activity was exhibited by NF1 0.00825 $\mu\text{M PNP g}^{-1}\text{soil min}^{-1}$ and lowest by D1 0.00219 $\mu\text{M PNP g}^{-1}\text{soil min}^{-1}$. All natural forest soil samples showed higher phosphatase activity than dump, which can be positively correlated with the dehydrogenase activity and negatively with catalase activity.




Graph 1: pH of different soil samples of Dump and Natural forest.




Graph 2: Comparison between colony forming units of natural forest and dump soil (all values in x10⁴ CFU/g)



Graph 5: Soil phosphatase activity of different Dump and Natural Forest soil sample (values in µM PNP g⁻¹ soil min.⁻¹).



Graph 3: Comparison between soil Catalase activity of Dump and Natural Forest Soil (all values in ml of .1 M KMnO₄ g⁻¹ of soil min.⁻¹).



Graph 4: Comparison of Soil Dehydrogenase activity between different samples of Dump and Natural forest (values expressed in µg TPF g⁻¹ dry soil.h⁻¹).

iv. Conclusion

This study was done to find the ill effects of mining over the environment. The microbial community and soil enzyme activity were taken as an indicator of environment, because these two parameters represent the health of the environment in different ways viz., microbes produce the soil enzymes and these enzymes regulate the biogeochemical cycles and these cycles play a vital role in maintaining the environment and ecosystem. The low number of microorganisms in the dump soil indicated towards contamination of heavy metals and stress conditions. The soil enzyme activity also co-related with the results as soil dehydrogenase and phosphatase activity has shown very low activity in the dump soil when compared to the Natural forest soil. The catalase activity confirmed the stress conditions in dump soils.

Acknowledgment

Authors are indebted to many who helped in field and laboratory during the course of the study. In addition, the authors would like to thank Department of Biotechnology, Govt. of India for providing financial assistance and Director, Motilal Nehru Institute of Technology Allahabad for extending research facilities.

In India, it has been estimated that about 10,000 hectares of land is affected by stockpiling of coal mine spoil. [30]

References

- [1] Johnson CD, Skousen JG (1995) Mine soil properties of 15 abandoned mine land sites in West Virginia. *J Environ Qual* 24: 635-643.
- [2] Jha AK, Singh JS (1991) Spoil characteristics and vegetation development of an age series of mine spoils in a dry tropical environment. *Vegetatio* 97: 63-76.
- [3] Mummey DL, Stahl PD, Buyer JS (2002) Soil microbiological properties 20 years after surface mine reclamation: spatial analysis of reclaimed and undisturbed sites. *Soil Biol Biochem* 34: 1717-1725.
- [4] Mc Sweeney K, Jansen IJ (1984) Soil structure and associated rooting behavior in mine soils. *Soil Sci Soc Am J* 48: 607-612.
- [5] Visser S, Zak J, Parkinson D (1979) Effects of surface mining on soil microbial communities and processes. Pregamon Press, NewYork, USA.
- [6] Juwarkar AA, Jambulkar HP, Singh SK (2004) Appropriate strategies for reclamation and revegetation of coal mine spoil dumps. Proceedings of the National Seminar on Environmental Engineering with special emphasis on Mining Environment, NSEEME-2004, India.
- [7] Ciolkosz EJ, Cronce RC, Cunningham RL, Peterson GW (1985) Characteristics, genesis, and classification of Pennsylvania mine soils. *Soil Science* 139: 232-238.
- [8] Roberts JA, Daniels WL, Bell JC, Burger JA (1988) Early stages of mine soil genesis as affected by top soiling and organic amendments. *Soil Sci Soc Am J* 52: 730-738.
- [9] Smejkalova, M., Mikanova, O., and Boruka, L. (2003) Effects of heavy metal concentrations on biological activity of soil microorganisms. *Plant Soil Environment*, 49(7), pp. 321-326
- [10] Nannipieri P, Greco S, Ceccanti B (1990) Ecological significance of the biological activity in soil. *Soil Biochem*. 6: 293-355.
- [11] Burns RG, (1982) Enzyme activity in soil: location and possible role in microbial ecology. *Soil Biology and Biochemistry*, 14: 423-427,
- [12] Allison S and Vitousek P, Responses of extracellular enzymes to simple and complex nutrient inputs. *Soil Biology and Biochemistry*, 37: 937-944, (2005).
- [13] Vangronsveld, J., and Cunningham, S. (1998) Introduction to the Concepts. In: Vangronsveld, J., and Cunningham, S, eds.. *Metal-Contaminated Soils: In Situ Inactivation and Phytoremediation*. Berlin/Heidelberg, Springer, pp. 1-15.
- [14] Renella, G., Mench, M., Gelsomin, A., Landi, L., and Nannipieri, P. (2005) Functional activity and microbial community structure in soils amended with bimetallic sludges. *Soil Biology and Biochemistry*, 37, pp. 1498-1506.
- [15] Akala B 1995 North Eastern coalfields – some highlights; *J. Mines, Metals and Fuels* 43(9-10) 303-305.
- [16] Chaoji V 2002 Environmental challenges and the future of Indian coal; *J. Mines, Metals and Fuels* 11 257-262.
- [17] Black, C.A., Evans, D.D., White, J.L., Ensminger, L.E. and Clark, F.E. (1965) *Methods of Soil Analysis: Chemical and Microbiological Properties*, Agronomy 9, Part II, Medison Wisconsin, USA.
- [18] Zhou L K, 1987. The Science of Soil Enzyme. The Science Press, Beijing, China. 267-270.
- [19] Von Mersi, W. and F. Schinner, 1991. An improved and accurate method for determining the dehydrogenase activity of soils with iodinitrotetrazolium chloride. *Biol. Fertil. Soils*, 11: 216-220.
- [20] Trevors JT (1984) Dehydrogenase activity in soil: A comparison between the INT and TTC assay. *Soil Biol Biochem* 16:673-674
- [21] Tabatabai MA, Bremner JM (1969) Use of p-nitrophenyl phosphate for assay of soil phosphatase activity. *Soil Biol Biochem* 1: 301-307.
- [22] Burger JA, Zipper CE (2002) How to restore forests on surface mined land. Powell river project, Virginia cooperative extension, USA 1-21.
- [23] Dutta RK, Agarwal M (2002) Effect of tree plantations on the soil characteristics and microbial activity of coal mine spoil land. *Trop Ecol* 43: 315-324.
- [24] Kandeler E, Dick R (2007) Soil enzymes: Spatial distribution and function in Agroecosystems. Taylor and Francis group, Boca Raton, usa.
- [25] G Alef K, Nannipieri P (1995) Methods in applied soil microbiology and biochemistry. Academic Press, London, UK.
- [26] Beyer L, Wackendorf C, Balzen FM, Balzer-Graf VR (1992) The effect of soil texture and soil management on microbial biomass and soil enzyme activities in arable soils of Northwest Germany. *Agrobiol Res* 45: 276.
- [27] Pascule JA, Hernanzed T, Garcia C, Ayuso M (1998) Enzymatic activities in an arid soil amended with urban organic wastes: laboratory experiment. *Bioresource Tech* 64: 131-138.
- [28] Caldwell BA (2005) Enzyme activities as a component of soil biodiversity: a review. *Pedobiologia* 49: 637-644.
- [29] Kramer S, Green DM (2000) Acid and alkaline phosphatase dynamics and their relationship to soil microclimate in a semi-arid woodland. *Soil Biol Biochem* 32:179-188.
- [30] Juwarkar, A.S., Thawale, P.R., Mowade, S., Shrivastav, S., Deshbhratar, P.B. and Juwarkar, A. (1994) 'Reclamation of coal mine spoil dump through integrated biotechnology approach', Proceedings of the International Symposium, VRCE Nagpur, India.

About Author (s):

Shobhit Verma & Abhay Pratap Singh: Students of Master of Technology in Biotechnology, at Department of Biotechnology, Motilal Nehru National Institute of Technology allahabad, Uttar Pradesh, India.

Shikha Devi: PhD research scholar in the Department of Biotechnology, Motilal Nehru National Institute of Technology allahabad, Uttar Pradesh, India.

Dr. Radha Rani Mewaram: Assistant Professor in the Department of Biotechnology, Motilal Nehru National Institute of Technology. Specialization in the field of Environmental Biotechnology and Phyto/bioremediation.

Dr. Shivesh Sharma: Head of the Department and Associate Professor in the Department of Biotechnology, Motilal Nehru National Institute of Technology Allahabad. Specialization in Environmental Biotechnology, Plant Microbe Interaction and Microbiology.

Dr. Kumud Dubey: Senior Scientist and Director at Centre for Social Forestry and Eco- rehabilitation Allahabad, India