

Production of laccase by *Resinicium bicolor* in submerged cultures: application of the Plackett-Burman experimental design to screen major factors

¹Harmen Shah, ²Faridah Yusof and ³Zahangir Alam

Abstract—The present work was aimed towards the evaluation of effects of the physical and chemical factors on laccase production by white rot fungus, *Resinicium bicolor* ATCC 64897 under submerged condition in shake flask using Plackett-Burman design. *R.bicolor* potentially detoxifies ground tire rubber (GTR) and facilitates growth of *Thiobacillus ferrooxidans*, a bacterium that performs devulcanization of GTR but is sensitive towards rubber additives present in GTR. In this experiment out of 11 factors screened, copper sulphate and yellow flame bark powder were found to be the most significant factors affecting the laccase production. The study indicated that the screened factors can be optimized for enhanced laccase production and can potentially be utilized for attaining maximum detoxification of GTR and hence further devulcanization.

Keywords— white-rot fungus, *Resinicium bicolor*, laccase, inducers

I. Introduction

Laccase (benzenediol: oxygen oxidoreductase, EC 1.10.3.2) belong to the copper metalloenzymes and to the blue oxidase sub-group that catalyzes reduction of molecular oxygen to water (1,2,3,4). White rot fungi (basidiomycetes) have a natural mechanism to decompose lignocellulosic materials and xenobiotics (5) with the help of their extracellular enzymes viz., lignin peroxidase, manganese peroxidase, versatile peroxidases, and laccases which allows them to decompose

Harmen Shah; Prof.Dr.Faridah Yusof, Prof.Dr.Zahangir Alam
International Islamic University Malaysia
Kuala Lumpur, Malaysia

Corresponding author.

organic matter esp. lignin and wood components and xenobiotic compounds (6,7,8). The production of laccase under submerged condition and the involvement of lignocellulosic materials and inducers in enhancing laccase production has been widely documented (9,10,11, 12,13,14,15,16,17). The ability of white rot fungus in synthetic dye decolorization and degradation of organo-pollutants and recalcitrant compounds have been extensively studied (18,19, 20, 21,22,23,24,25).

The devulcanization potential of *Thiobacillus ferrooxidans* has been well proven by several research groups (26,27,28,29) but in most cases *T. ferrooxidans* was found sensitive towards additives like accelerants, stabilizers etc added during rubber vulcanization process (30,31). In a study it was demonstrated that the microbial toxicity towards rubber constituents have profound effect on devulcanization process whereby a white rot fungus *Resinicium bicolor* has been proven to be very effective in detoxifying ground tyre rubber under submerged culture condition with its extracellular ligninolytic enzymes (32). Thus the objective of this study was to screen out the major factors affecting laccase production in *R. bicolor* under submerged culture using plackett-burman design (PBD). Thus, this work is an initiative towards the detoxification of ground tire rubber using the laccase mediator system (LMS) of *R. bicolor* to facilitate the devulcanization process by bacteria *T. ferrooxidans*.

II Materials and Methods

A. Fungal strain

The fungus *Resinicium bicolor* strain 64897 used in this study was obtained from ATCC (American Type Culture Collection, Manassas, USA). Stock cultures were grown at room temperature (26-28°C) for 3-4 days, maintained on malt agar (ATCC® Medium 323) plates and stored at 4°C with subsequent transfers every 4-5 weeks.

B. Substrate

The lignocellulosic substrate consisted Cassava peel, Yellow flower tree bark and Rubber tree bark. These were crushed in a mechanical crusher (HITOP-SY-20) and ground to 0.5 mm in a rotomill- Pulverisette 14 (Fritsch, Germany) and sieved through 200 mesh to get powder with uniform particle size to be used as carbon source in fermentation medium.

C. Chemicals

p-Coumaric acid, 1-Hydroxybenzotriazole, Syringaldazine and 2,5-Xylidine of analytical grade were purchased from Sigma-Aldrich (St. Louis, MO). Copper sulphate and buffer salts were from HmbG (Germany). Malt extract (LP0039) was from Oxoid (England) and malt extract broth was from Merck Darmstadt (Germany). Bacto™ Agar was from Difco (USA).

D. Fermentation medium

The malt extract broth 1.7% (w/v) was used both as nitrogen source and fermentation medium. The fermentation medium was supplemented by inducers and substrates in the concentration as per values from plackett-burman design (PBD) and was autoclaved at 1.5 atmosphere and 121°C for 15 minutes.

E. Inoculum preparation and submerged cultivation

Inocula were prepared as spore suspensions by scraping a 4 days old malt agar slant with 20 ml sterile distilled water. Each 250 ml shaker flask containing 100 ml culture medium received 10 ml of this inoculum and pH was adjusted using 1M HCl. All incubations were performed in an incubator shaker in the dark with three replicates under conditions according to PBD.

F. Plate assay

To determine the presence of extracellular laccase, a plate assay using method of (33) was performed. The petri plates containing 20 ml of 4.5% malt agar was supplemented with 0.08% (w/v) of bromophenol blue. The fungal discs of 7 mm were incised from the periphery of actively growing plate of *R. bicolor* and were aseptically transferred to plates containing bromophenol blue. The plates were incubated for 1 day at room temperature. The formation of clear halo around mycelia was considered to be a positive test for extracellular laccase.

G. Estimation of laccase activity

Laccase assay was performed by using the method of (34). The reaction mixture contained 0.3 ml of 0.216 mM syringaldazine, 2.2 ml of 100 mM citrate- phosphate buffer; pH 6.5 (35) and 0.5 ml of crude enzyme sample obtained by centrifuging culture broth at 10,000 rpm for 10 minutes. Oxidation of syringaldazine was followed by increase in absorbance at 530 nm. Enzyme activity was expressed in units/ml (U/ml). One unit produced a ΔA_{530} of 0.001 per minute at pH 6.5 at 30°C in a 3 ml reaction volume using syringaldazine as substrate. The change in absorbance was recorded by a spectrophotometer (ViVa spec. LS, Sartorius stedim, Germany).

H. Screening using Plackett-Burman design

The Plackett-Burman method is an effective two-level factorial design, which identifies the crucial factors required for enhanced enzyme production by screening N variables in N+1 experiments (36) whereby interaction between various factors is not taken into account. The screened variables (factors) that are significant and influential can further be

optimized using response surface methodology. To determine the major factors significantly influencing the laccase activity in *Resinicium bicolor* strain ATCC 64897, a set of 12 runs were conducted using 11 independent variables or factors that included 8 chemical factors and 3 physical factors wherein each factor was performed at two levels i.e; low and high as shown in table I. Each run was performed in triplicate and mean value and standard deviation (Mean \pm S.D) of response (laccase activity) was measured.

The complete experimental design with different combination of factors in each run has been depicted in table II.

TABLE I. LEVELS OF THE VARIABLES IN PLACKETT-BURMAN DESIGN

Variables	Level		
	Factors	Low	High
Chemical factor (X ₁)	Copper sulphate (mM)	0	5
Chemical factor (X ₂)	p-Coumaric Acid (mM)	0	1
Chemical factor (X ₃)	1-Hydroxybenzotriazole (mM)	0	1
Chemical factor (X ₄)	2,5-Xylidine (mM)	0	1
Chemical factor (X ₅)	Temperature (°C)	24	28
Chemical factor (X ₆)	Cassava peel powder (g)	1	5
Chemical factor (X ₇)	Yellow flower tree bark powder (g)	1	5
Chemical factor (X ₈)	Rubber tree bark powder (g)	1	5
Physical factor (X ₉)	Agitation (rpm)	140	180
Physical factor (X ₁₀)	pH	2.5	5.5
Physical factor (X ₁₁)	Incubation period (days)	1	4

TABLE II. PLACKETT-BURMAN DESIGN FOR LACCASE PRODUCTION

Run	X ₁	X ₂	X ₃	X ₄	X ₅	X ₆	X ₇	X ₈	X ₉	X ₁₀	X ₁₁
1	5	0	0	0	28	5	5	1	180	5.5	1
2	0	0	1	1	28	1	5	5	140	5.5	1
3	0	0	0	0	24	1	1	1	140	2.5	1
4	0	1	0	0	24	5	5	5	140	5.5	4
5	5	0	1	0	24	1	5	5	180	2.5	4

6	5	0	0	0	28	1	1	1	180	5.5	1
7	0	0	1	1	28	1	1	1	140	5.5	1
8	0	0	0	0	24	1	1	1	140	2.5	1
9	0	1	0	0	24	5	5	5	140	5.5	4
10	5	0	1	0	24	1	5	5	180	2.5	4
11	0	0	0	1	28	5	1	5	180	2.5	4
12	0	1	1	0	28	1	1	1	180	5.5	4

Proc. of the Intl. Conf. on Advances in Civil, Structural, Environmental & Bio-Technology -- CSEB 2014
 Copyright © Institute of Research Engineers and Doctors. All rights reserved.
 ISBN: 978-1-63248-001-9 doi: 10.15224/ 978-1-63248-001-9-105

III Results and Discussion

A. Plate assay

Plate assay is rapid technique of identifying the enzyme of concern in a particular strain and is reported by several groups using substrates like syringaldazine, guaiacol and 2, 2'-azino-bis (3-ethylbenzthiazoline-6-sulfonic acid) (ABTS) (37,38,39,40). The strain *R. bicolor* gave positive reaction in plate assay using the azo dye, bromophenol blue as substrate for extracellular laccase which was visualized by the appearance of halo around mycelia as shown in figure 1. The halo appeared after 24 hours of incubation in the absence of any mediators which shows that the laccase secreted by *R. bicolor* can mediate the decolorization of synthetic dyes even in absence of any mediator in a short period as decolorization of azo dyes were performed in the presence of mediators (41).




Figure 1. Plate assay method for laccase test showed the yellowish halo developed by *R. bicolor* using bromophenol blue as substrate after 24 hours of incubation.

B. Screening of factors using plackett-burman design

To screen out the major factors responsible for increased laccase production, 11 factors comprising chemical inducers, lignocellulosic materials, agitation, pH and incubation period were compared at two levels. The results of screening experiment has been presented in figure 2. On the basis of these results, the main effect of each factor on laccase activity was

calculated using (1). The contribution of different factors towards main effect has been depicted in figure 3. A broad variation in the laccase activity ranging from 0.53 to 6.8 (U/ml) was found. This variation is an indication for optimization of screened factors for enhanced laccase productivity.

$$\text{Main effect} = \frac{\Sigma \text{Response value from high level} - \Sigma \text{Response value from low level}}{\Sigma \text{Number of runs}}$$




Figure 2. Laccase activity as response for plackett-burman screening.




Figure 3. Different factors and their contribution towards main effect.

As from figure 3, it is clear that the activity of laccase in case of *R. bicolor* is primarily influenced by three major factors which were copper sulphate (X₁), yellow flame bark (X₇) and incubation period (X₁₁). 2,5-xylidine (X₄) and pH (X₁₀) have little positive effect on laccase activity and p-coumaric acid (X₂) and cassava peel (X₆) gave negative effect on laccase activity while effect of 1-hydroxybenzotriazole (X₃) and rubber bark powder (X₈) were negligible. Thus X₁, X₅, X₇, X₉ and X₁₁ were selected for analysis. The analysis of variance of experimental design showed that X₁, X₇ and X₁₁ are significant model terms (table III). The final regression equation developed by plackett-burman design in terms of coded factors was as follows:

$$R_1 = 3.04 + 1.30X_1 + 0.39X_5 + 1.15X_7 + 0.20X_9 - 0.56X_{11} \quad (2)$$

Where 'R₁' is the laccase enzyme activity (U/ml) and X₁, X₅, X₇, X₉ and X₁₁ are coded factors.

TABLE III. Analysis of variance (ANOVA) for selected

	Sum of		Mean	F		
Source	Squares	DF	Square	Value	Prob > F	
Model	42.18	5	8.44	16.09	0.0020	significant
X ₁	20.13	1	20.13	38.41	0.0008	significant
X ₅	1.86	1	1.86	3.55	0.1087	
X ₇	15.89	1	15.89	30.32	0.0015	significant
X ₉	0.50	1	0.50	0.96	0.3652	
X ₁₁	3.79	1	3.79	7.23	0.0361	significant
Residual	3.15	6	0.52			
Cor Total	45.32	11				
Std. Dev.	0.72				R ²	0.9306
Mean	3.04				Adj R ²	0.8728
C.V.	23.83				Pred R ²	0.7224
PRESS	12.58				Adeq Precision	11.8932

factorial model.

The Model F-value of 16.09 implied that the model is significant. Values of "Prob>F" and less than 0.05 indicated that model terms are significant. In this case copper sulphate (X₁), yellow flame bark powder (X₇) and incubation period (X₁₁) were considered as significant model terms (factors) with confidence level greater than 95% (P<0.05). The "Predicted R-Squared" of 0.7224 is in reasonable agreement with the "Adjusted R-Squared" of 0.8728. The coefficient of determination (R²) of the model was calculated to be 0.9306, which indicated that 93.06% of the variability in the response could be explained by the model.

C. Copper/lignocellulose synergistic induction of laccase

The inducers employed in this experiment consisted of copper sulphate, p-coumaric acid, 1-hydroxybenzotriazole and 2,5-xylidine whereas the lignocellulosic materials used in the current study were: cassava peel powder, yellow flame tree bark powder and rubber tree bark powder. Out of these two categories, the copper sulphate and yellow flame bark powder was found to be significant in enhancing the laccase activity. The highest laccase activity was 6.80 U/ml. The increment in the laccase production using lignocellulosic materials and copper sulphate as inducer has already been established by several research groups including (42, 43). To our knowledge this is the first attempt to determine the effect of Yellow flame bark powder as lignocellulosic material for laccase production in strain *R. bicolor*.

The lignocellulosic material being as rich carbohydrate source increases laccase activity (44) and the major portion of total phenolic content of yellow flame bark consist of tannins (45) especially tannic acid which has been shown to act as a good substrate as well as an inducer for laccase production in different white rot fungi (46, 47).

D. Incubation period and its effect on laccase production

The strain exhibited laccase activity at the end of first day of incubation. Thus, it can be said that availability of fresh culture

medium supplemented with lignocellulosic substrates and inducers under agitation condition supported the growth and metabolic reaction in fungal species which resulted in laccase production.

Conclusion

The white rot fungal strain *R. bicolor* has the ability to produce laccase under submerged culture condition using cheap substrates and inducers. The placket-burman screening showed three significant factors affecting laccase production viz., copper sulphate, yellow flame bark powder and incubation period which can be further optimized to get maximum laccase production and can be utilized for optimum detoxification ground tire rubber. Till date there is no data available for the detection of laccase in *R. bicolor*.

Acknowledgment

This work was conducted using the e-science fund (SF-020-0049) granted by Ministry of Sciences, Technology and Innovation, Malaysia.

References

- 1) P. Baldrian, "Fungal laccases: occurrence and properties," *FEMS Microbiol Rev.* 30, 215–242, (2005).
- 2) O.V. Morozova, G.P. Shumakovich, M.A. Gorbacheva, S.V. Shleev, and A.I. Yaropolov, "Blue Laccases" *Biochemistry (Moscow)*. 72(10), 1136-1150, (2007).
- 3) C.J. Rodgers, C.F. Blanford, S.R. Giddens, P. Skamnioti, F.A. Armstrong, and S.J. Gurr, "Designer laccases: a vogue for high-potential fungal enzymes," *Trends in Biotechnology*. 28 (2), 63-72, (2009).
- 4) Shradha, R. Shekher, S. Sehgal, M. Kamthania, and A. Kumar, "Laccase: Microbial Sources, Production, Purification, and Potential Biotechnological Applications," *Enzyme Research*, Article ID 217861, 11 pages, (2011).
- 5) N.A. Kulikova, O.I. Klein, E.V. Stepanova and O.V. Koroleva, "Use of Basidiomycetes in Industrial Waste Processing and Utilization Technologies: Fundamental and Applied Aspects (Review)" *Applied Biochemistry and Microbiology*. 47(6), 565–579, (2011).
- 6) T.K. Kirk and R.L. Farrell, "Enzymatic "combustion": the microbial degradation of lignin" *Ann Rev Microbiol*. 41, 465–505, (1987).
- 7) Ward, Y. Hadar, and C.G. Dosoretz, "The Biodegradation of Lignocellulose by White Rot Fungi," 33, Marcel Dekker, Inc. (2004).
- 8) A. Kunamneni, F.J. Plou, A. Ballesteros, and M. Alcalde, "Laccases and Their Applications: A Patent Review" *Recent Patents on Biotechnology*. 2, 10-24, (2008).
- 9) M. Lorenzo, D. Moldes, S. Rodriguez-Couto, and K. Sanroman, "Improving laccase production by employing different lignocellulosic wastes in submerged cultures of *Trametes versicolor*" *Bioresource technology*. 82, 109-113, (2002).
- 10) D. Moldes, M. Lorenzo, and M.A. Sanroman, "Different proportions of laccase isoenzymes produced by submerged cultures of *Trametes versicolor* grown on lignocellulosic wastes" *Biotechnology Letters*. 26, 327–330, (2004).
- 11) M.S. Revankar, "Enhanced production of laccase using a new isolate of white rot fungus WR-1," *Process Biochemistry*. 41(3), 581-588, (2006).
- 12) E. Birhanli, and O. Yesilada, "Enhanced production of laccase in repeated-batch cultures of *Funalia trogii* and *Trametes versicolor*," *Biochemical Engineering Journal*. 52, 33–37, (2010).
- 13) R. Tinoco, A. Acevedo, E. Galindo, and L. Serrano-Carreón, "Increasing *Pleurotus ostreatus* laccase production by culture medium optimization and copper/lignin synergistic induction," *J Ind Microbiol Biotechnol*. 38, 531–540, (2011).

- 14) J.D. Lebrun, I. Lamy, and C. Mougin, "Favouring the bioavailability of Zn and Cu to enhance the production of lignin-modifying enzymes in *Trametes versicolor* cultures" *Bioresource Technology*. 102, 3103–3109, (2011).
- 15) G.S. Dhillon, S. Kaur and S.K. Brar, " In-vitro decolorization of recalcitrant dyes through an ecofriendly approach using laccase from *Trametes versicolor* grown on brewer's spent grain" *International Biodeterioration and Biodegradation*. 72, 67-75, (2012).
- 16) K. Fu, S. Fu, H. Zhan, P. Zhou, M. Liu, and H. Liu, "A newly isolated wood-rot fungus for laccase production in submerged cultures" *BioResources*, 8(1), 1385-1397, (2013).
- 17) Z. Ding, Y. Chen, Z. Xu, L. Peng, G. Xu, Z. Gu, L. Zhang, G. Shi, and K. Zhang, " Production and characterization of laccase from *Pleurotus ferulae* in submerged fermentation" *Ann Microbiol.* (2013).
- 18) K. Selvam, K. Swaminathan, and K.S. Chae, "Decolourization of azo dyes and a dye industry effluent by a white rot fungus *Telephora* sp.," *Bioresource Technology*. 88, 115–119, (2003).
- 19) G. Eibes, T. Cajthaml, M. Moreira, G. Feijoo and J. Lema, "Enzymatic degradation of anthracene, dibenzothiophene and pyrene by manganese peroxidase in media containing acetone" *Chemosphere*. 64, 408–414, (2006).
- 20) I. Canas, M. Alcalde, F. Plou, M. Martiane, A. Martiane, & S. Camarero, "Transformation of Polycyclic Aromatic Hydrocarbons by Laccase Is Strongly Enhanced by Phenolic Compounds Present in Soil" *Environ. Sci. Technol.* 41, 2964-2971, (2007).
- 21) K. Svobodova, A. Majcherczyk, A. Novotny, and U. Kues, " Implication of mycelium-associated laccase from *Irpex lacteus* in the decolorization of synthetic dyes," *Bioresource Technology*. 99, 463–471, (2008).
- 22) L. Singh and V.P. Singh, "Biodegradation of Textile Dyes, Bromophenol blue and Congo red by Fungus *Aspergillus Flavus*," *Environ. We Int. J. Sci. Tech.* 5, 235-242, (2010).
- 23) M. Neifar, A. Kamoun, A. Jaouani, R. Ellouze-Ghorbel, and S. Ellouze-Chaabouni, " Application of Asymmetrical and Hoke Designs for Optimization of Laccase production by the White-Rot Fungus *Fomes fomentarius* in solid-State fermentation," *Enzyme Research*. Article ID 368525, 12 pages, (2011).
- 24) M.E. Urrea, C.A. Reddy, and S.N. Singh, Chapter, 2: Microbial Degradation of Xenobiotics. (ed.), (2012). *Environmental Science and Engineering*. DOI: 10.1007/978-3-642-23789-8_2, Springer-Verlag Berlin Heidelberg
- 25) D. Daassi, H. Zouari-Mechichi, F. Frikha, M.J. Martinez, M. Nasri and T. Mechichi, "Decolorization of the azo dye Acid Orange 51 by laccase produced in solid culture of a newly isolated *Trametes trogii* strain" *3 Biotech*. 3, 115–125, (2013).
- 26) Torma A.E and Raghavan D, "Biodesulphurization of rubber materials" In: *Bioprocess Engineering Symposium*, American Society of Mechanical Engineers, Bioengineering Division. 16, 81-87, (1990).
- 27) M. Löffler, G. Straube, and E. Straube, "Biohydrometallurgical Technologies: Proceedings of an International Biohydrometallurgy Symposium, presented at Biohydrometallurgical Symposium" *Jackson hole, Wyo.* 2, 673, (1993).
- 28) R. Romine, L. Snowden, and L. Swan, "Method for the addition of vulcanised waste rubber to virgin rubber products," *US/5,597,851*; (1997).
- 29) Y. Li, S. Zhao, and Y. Wang, " Microbial desulphurization of ground tire rubber by *Thiobacillus ferrooxidans*" *Polymer Degradation and Stability*. 96, 1662-1668, (2011).
- 30) B.J. Zyska, "Microbial deterioration of rubber," *Biodeterioration*. 7, 535-552, (1988).
- 31) M. Christiansson, B. Stenberg, and O. Holst, "Toxic additives- a problem for microbial waste rubber desulfurization" *Resource Environ Biotechnol* 3. 11–21,(2000).
- 32) K. Bredberg, K. Andersson, E. Landfors, and O. Holst, "Microbial detoxification of waste rubber material by wood-rotting fungi," *Biosource technology*. 83, 221-224, (2002).
- 33) Tekere; Growth, dye degradation and ligninolytic activity studies on Zimbabwean white rot fungi," *Enzyme and Microbial Technology*; 28, 420–426, (2001).
- 34) J.P. Ride, *Physiological Plant Pathology*. 16: 187-196, (1980).S.R Couto and J.L.T Herrera, "Laccase production at reactor scale by filamentous fungi," *Biotechnol Adv.* 25, 6, 500-513 (2007).
- 35) G. Gomori, *Preparation of Buffers for Use in Enzyme Studies. In Methods in Enzymology*, Vol. II, Colowick, S. P., Kaplan, N. O. (Eds.); Academic Press: New York, p. 138-146, (1955).
- 36) R.L. Plackett and J.P. Burman, " The design of optimum multifactorial experiments," *Biometrika*, 33(4), 305–325, 1946
- 37) L.L. Kiiskinen, M. Ratto, and K. Kruus, "Screening for novel laccase-producing microbes" *Journal of Applied Microbiology*. 97, 640–646, (2004).
- 38) V.V. Kumar, S.D. Kirupha, P. Periyaraman, and S. Sivanesan, "Screening and induction of laccase activity in fungal species and its application in dye decolorization" *African Journal of Microbiology Research* Vol. 5(11), pp. 1261-1267, 4 June (2011).
- 39) S.S. More, P. S. Renuka, K. Pruthvi, M. Swetha, S. Malini, and S. M. Veena, "Isolation, Purification, and Characterization of Fungal Laccase from *Pleurotus* sp." *Enzyme Research*. Article ID 248735, 7 pages (2011).
- 40) F. Sheikhi, M.A. Roayaei, N. Enayatizamir, and S. Rodriguez-Couto, "The Determination of Assay for Laccase of *Bacillus subtilis* WPI with Two Classes of Chemical Compounds as Substrates," *Indian J Microbiol.* 52(4),701–707, Oct–Dec (2012).
- 41) S. Camarero, D. Ibarra, M.J. Martinez, and A.T. Martinez, "Lignin-Derived Compounds as Efficient Laccase Mediators for Decolorization of Different Types of Recalcitrant Dyes" *Applied and Environmental Microbiology*, Vol. 71, No. 4p. 1775–1784, Apr. (2005).
- 42) S.S. Kahraman and I.H. Gurdal, "Effect of synthetic and natural culture media on laccase production by white rot fungi", *Bioresource Technology* 82, 215–217, (2002).
- 43) K.N. Niladevi, R.K. Sukumaran, N. Jacob, G.S. Anisha, and P. Prema, "Optimization of laccase production from a novel strain-*Streptomyces psammoticus* using response surface methodology," *Microbiological Research*. 164, 105-113, (2009).
- 44) S. Rodriguez Couto and M.A. Sanroman, "Coconut flesh: a novel raw material for laccase production by *Trametes hirsuta* under solid state conditions: Application to Lissamine Green B decolourization," *J Food Eng.* 71, 208–13, (2005).
- 45) T. Manaharan, L.L. Teng, D. Appleton, C.H. Ming, T. Masilamani, and U.D. Palanisamy, "Antioxidant and antiglycemic potential of *Peltophorum pterocarpum* plant parts," *Food Chemistry*. 129, 1355–1361, (2011).
- 46) R. Bourbonnais, M.G. Paice, B. Freiermuth, E. Bodie, and S. Borneman, "Reactivities of various mediators and laccases with kraft pulp and lignin model compounds," *Applied and Environmental Microbiology*, vol. 63, no. 12, pp. 4627–4632, (1997).
- 47) H.Y. Alfara, N.H.M. Hasali, and Omar M.N, "A lignolytic Fungi with Laccase Activity Isolated from Malaysian local Environment for Phytochemical Transformation Purposes" *Int. Res. J. Biological Sci.* Vol. 2(2), 51-54, February (2013).