

# On the Existence of $n$ -Tuple Magic Rectangles

Phaisatcha Inpoonjai and Thiradet Jiarasuksakun

**Abstract**—Magic rectangles are a classical generalization of the well-known magic squares, and they are related to graphs. A graph  $G$  is called degree-magic if there is a labelling of the edges by integers  $1, 2, \dots, |E(G)|$  such that the sum of the labels of the edges incident with any vertex  $v$  is equal to  $(1+|E(G)|)\deg(v)/2$ . In this paper we generalize magic rectangles to be  $n$ -tuple magic rectangles, and prove the necessary and sufficient conditions for the existence of even  $n$ -tuple magic rectangles. Using this existence we identify the sufficient condition for degree-magic labellings of the  $n$ -fold self-union of complete bipartite graphs to exist.

**Keywords**—magic squares, magic rectangles, degree-magic graphs

## I. Introduction

Magic rectangles are a natural generalization of the magic squares which have widely intrigued mathematicians and the general public. A magic  $(p, q)$ -rectangle  $R$  is a  $p \times q$  array in which the first  $pq$  positive integers are placed such that the sum over each row of  $R$  is constant and the sum over each column of  $R$  is another (different if  $p \neq q$ ) constant. Harmuth [1, 2] studied magic rectangles over a century ago and proved that

**Theorem 1** ([1, 2]) *For  $p, q > 1$ , there is a magic  $(p, q)$ -rectangle  $R$  if and only if  $p \equiv q \pmod{2}$  and  $(p, q) \neq (2, 2)$ .*

In 1990, Sun [3] studied the existence of magic rectangles. Later, Bier and Rogers [4] studied balanced magic rectangles, and Bier and Kleinschmidt [5] studied centrally symmetric and magic rectangles. Then Hagedorn [6] presented a simplified modern proof of the necessary and sufficient conditions for a magic rectangle to exist. The concept of magic rectangles was generalized to  $n$ -dimensions and several existence theorems were proven by Hagedorn [7].

For simple graphs without isolated vertices, if  $G$  is a graph, then  $V(G)$  and  $E(G)$  stand for the vertex set and the edge set of  $G$ , respectively. Cardinalities of these sets are called the *order* and *size* of  $G$ .

Let a graph  $G$  and a mapping  $f$  from  $E(G)$  into positive integers be given. The *index mapping* of  $f$  is the mapping  $f^*$  from  $V(G)$  into positive integers defined by

$$f^*(v) = \sum_{e \in E(G)} \eta(v, e)f(e) \quad \text{for every } v \in V(G), \quad (1)$$

where  $\eta(v, e)$  is equal to 1 when  $e$  is an edge incident with a vertex  $v$ , and 0 otherwise. An injective mapping  $f$  from  $E(G)$  into positive integers is called a *magic labelling* of  $G$  for an *index*  $\lambda$  if its index mapping  $f^*$  satisfies

$$f^*(v) = \lambda \quad \text{for all } v \in V(G). \quad (2)$$

A magic labelling  $f$  of a graph  $G$  is called a *supermagic labelling* if the set  $\{f(e) : e \in E(G)\}$  consists of consecutive positive integers. A graph  $G$  is *supermagic (magic)* whenever a supermagic (magic) labelling of  $G$  exists.

A bijective mapping  $f$  from  $E(G)$  into  $\{1, 2, \dots, |E(G)|\}$  is called a *degree-magic labelling* (or only *d-magic labelling*) of a graph  $G$  if its index mapping  $f^*$  satisfies

$$f^*(v) = \frac{1+|E(G)|}{2} \deg(v) \quad \text{for all } v \in V(G). \quad (3)$$

A d-magic labelling  $f$  of  $G$  is called *balanced* if for all  $v \in V(G)$ , the following equation is satisfied

$$\left| \left\{ e \in E(G) : \eta(v, e) = 1, f(e) \leq \lfloor |E(G)|/2 \rfloor \right\} \right| = \left| \left\{ e \in E(G) : \eta(v, e) = 1, f(e) > \lfloor |E(G)|/2 \rfloor \right\} \right|. \quad (4)$$

A graph  $G$  is *degree-magic (balanced degree-magic)* or only *d-magic* when a d-magic (balanced d-magic) labelling of  $G$  exists.

The concept of magic graphs was introduced by Sedláček [8]. Later, supermagic graphs were introduced by Stewart [9]. There are now many papers published on magic and supermagic graphs; we refer the reader to Gallian [10] for more comprehensive references. Recently, the concept of degree-magic graphs was introduced by Bezegová and Ivančo [11] as an extension of supermagic regular graphs. They also established the basic properties of degree-magic graphs and proved that

**Proposition 1** ([11]) *For  $p, q > 1$ , the complete bipartite graph  $K_{p,q}$  is d-magic if and only if  $p \equiv q \pmod{2}$  and  $(p, q) \neq (2, 2)$ .*

**Theorem 2** ([11]) *The complete bipartite graph  $K_{p,q}$  is balanced d-magic if and only if the following statements hold:*

- (i)  $p \equiv q \equiv 0 \pmod{2}$ ;
- (ii) if  $p \equiv q \equiv 2 \pmod{4}$ , then  $\min\{p, q\} \geq 6$ .

In this paper we introduce  $n$ -tuple magic rectangles. To show their existence, we introduce the closely related concept

---

Phaisatcha Inpoonjai<sup>1</sup> and Thiradet Jiarasuksakun<sup>2</sup>  
 Department of Mathematics, Faculty of Science, King Mongkut's University of Technology Thonburi  
 126 Pracha Uthit Rd., Bang Mod, Thung Khru, Bangkok 10140, Thailand

of centrally  $n$ -tuple symmetric rectangles. Then we use the existence of centrally  $n$ -tuple symmetric rectangles to give a construction of even  $n$ -tuple magic rectangles. Finally, we identify the sufficient condition for  $d$ -magic labellings of the  $n$ -fold self-union of complete bipartite graphs to exist.

## II. The $n$ -Tuple Magic Rectangles

In this section we introduce  $n$ -tuple magic rectangles and prove the necessary and sufficient conditions for even  $n$ -tuple magic rectangles to exist.

**Definition 1** An  $n$ -tuple magic  $(p, q)$ -rectangle  $R := (r_{i,j}^1)(r_{i,j}^2) \dots (r_{i,j}^n)$  is a class of  $n$  arrays in which each array has  $p$  rows and  $q$  columns, and the first  $npq$  positive integers are placed such that the sum over each row of any array of  $R$  is constant and the sum over each column of  $R$  is another (different if  $p \neq q$ ) constant.

Let  $R := (r_{i,j}^1)(r_{i,j}^2) \dots (r_{i,j}^n)$  be an  $n$ -tuple magic  $(p, q)$ -rectangle. As each row sum of any array of  $R$  is  $q(npq+1)/2$  and each column sum of  $R$  is  $p(npq+1)/2$  and both are integer, we then have

**Proposition 2** If  $R$  is an  $n$ -tuple magic  $(p, q)$ -rectangle, then the following statements hold:

- (i) if  $n$  is odd, then  $p \equiv q \pmod{2}$ ;
- (ii) if  $n$  is even, then  $p \equiv q \equiv 0 \pmod{2}$ .

Proposition 2 allows the set of  $n$ -tuple magic rectangles to be divided into sets of odd and even rectangles. We quickly see that an  $n$ -tuple magic  $(2, 2)$ -rectangle does not exist. To show the existence of other even  $n$ -tuple magic rectangles, we introduce the closely related concept of centrally  $n$ -tuple symmetric  $(p, q)$ -rectangles as follows.

**Definition 2** Let  $x > -1$  and let  $R$  be a class of  $n$  even rectangular arrays in which each array has  $p$  rows and  $q$  columns and the entries of  $R$  are numbers  $\pm(x+1), \dots, \pm(x+npq/2)$ .  $R$  is a centrally  $n$ -tuple symmetric  $(p, q)$ -rectangle of type  $x$  if the sum over each row and column of any array is zero. Additionally, if  $R$  has an equal number of positive and negative numbers in each row and column of any array, we say that  $R$  is balanced.

If  $R$  is an even  $n$ -tuple magic  $(p, q)$ -rectangle, then by subtracting  $(npq+1)/2$  from each entry of  $R$ , we obtain a centrally  $n$ -tuple symmetric  $(p, q)$ -rectangle of type  $-1/2$ . Similarly, every centrally  $n$ -tuple symmetric  $(p, q)$ -rectangle of type  $-1/2$  determines an even  $n$ -tuple magic  $(p, q)$ -rectangle. Thus, we can use the existence of centrally  $n$ -tuple symmetric  $(p, q)$ -rectangles to prove the existence of even  $n$ -tuple magic  $(p, q)$ -rectangles.

**Lemma 1** For  $x, y > -1$ , if a balanced centrally  $n$ -tuple symmetric  $(p, q)$ -rectangle of type  $x$  exists, then a balanced centrally  $n$ -tuple symmetric  $(p, q)$ -rectangle of type  $y$  exists.

**Proof.** Suppose that  $R := (r_{i,j}^1)(r_{i,j}^2) \dots (r_{i,j}^n)$  is the given rectangle. Then we define a  $(p, q)$ -rectangle  $S := (s_{i,j}^1)(s_{i,j}^2) \dots (s_{i,j}^n)$  by

$$s_{i,j}^t = (y-x)\text{sgn}(r_{i,j}^t) + r_{i,j}^t, \quad \text{for every } t \in \{1, 2, \dots, n\}.$$

The entries of  $S$  are the numbers  $\pm(y+1), \dots, \pm(y+npq/2)$ . For any  $t \in \{1, 2, \dots, n\}$  and  $1 \leq i \leq p$ , the sum of each row is

$$\begin{aligned} \sum_{j=1}^q s_{i,j}^t &= \sum_{j=1}^q ((y-x)\text{sgn}(r_{i,j}^t) + r_{i,j}^t) \\ &= (y-x) \sum_{j=1}^q \text{sgn}(r_{i,j}^t) + \sum_{j=1}^q r_{i,j}^t = 0, \end{aligned}$$

and for all  $1 \leq j \leq q$ , the sum of each column is

$$\begin{aligned} \sum_{i=1}^p s_{i,j}^t &= \sum_{i=1}^p ((y-x)\text{sgn}(r_{i,j}^t) + r_{i,j}^t) \\ &= (y-x) \sum_{i=1}^p \text{sgn}(r_{i,j}^t) + \sum_{i=1}^p r_{i,j}^t = 0. \end{aligned}$$

Thus,  $S$  is a centrally  $n$ -tuple symmetric  $(p, q)$ -rectangle of type  $y$ . For any  $t \in \{1, 2, \dots, n\}$ , if  $r_{i,j}^t$  is positive, then  $r_{i,j}^t = x+m$  for some  $m \geq 1$ . Hence,  $s_{i,j}^t = y+m$  is also positive. Similarly,  $r_{i,j}^t$  negative implies  $s_{i,j}^t$  negative. Therefore,  $S$  is balanced.  $\square$

**Proposition 3** If a balanced centrally  $n$ -tuple symmetric  $(p, q)$ -rectangle exists, then an  $n$ -tuple magic  $(p, q)$ -rectangle exists.

**Proof.** Suppose  $R$  is the given rectangle. If  $R$  has type  $x$ , then by Lemma 1, there exists a balanced centrally  $n$ -tuple symmetric  $(p, q)$ -rectangle of type  $-1/2$ . Therefore, an  $n$ -tuple magic  $(p, q)$ -rectangle exists.  $\square$

**Example 1** We consider a balanced centrally 5-tuple symmetric  $(4, 2)$ -rectangle  $R := (r_{i,j}^1)(r_{i,j}^2) \dots (r_{i,j}^5)$  of type 1 as follows.

$$R := \begin{pmatrix} 2 & -2 \\ -3 & 3 \\ -4 & 4 \\ 5 & -5 \end{pmatrix} \begin{pmatrix} 6 & -6 \\ -7 & 7 \\ -8 & 8 \\ 9 & -9 \end{pmatrix} \begin{pmatrix} 10 & -10 \\ -11 & 11 \\ -12 & 12 \\ 13 & -13 \end{pmatrix} \begin{pmatrix} 14 & -14 \\ -15 & 15 \\ -16 & 16 \\ 17 & -17 \end{pmatrix} \begin{pmatrix} 18 & -18 \\ -19 & 19 \\ -20 & 20 \\ 21 & -21 \end{pmatrix}.$$

Then we define a 5-tuple  $(4, 2)$ -rectangle  $S := (s_{i,j}^1)(s_{i,j}^2) \dots (s_{i,j}^5)$  related to  $R$  by

$$s_{i,j}^t = -\frac{3}{2}\text{sgn}(r_{i,j}^t) + r_{i,j}^t, \quad \text{for every } t \in \{1, 2, 3, 4, 5\}.$$

Thus,  $S$  is a balanced centrally 5 -tuple symmetric  $(4, 2)$  -rectangle  $S$  of type  $-1/2$  as follows.

$$S := \begin{pmatrix} \frac{1}{2} & -\frac{1}{2} \\ \frac{3}{2} & \frac{3}{2} \\ -\frac{5}{2} & \frac{5}{2} \\ \frac{7}{2} & -\frac{7}{2} \end{pmatrix} \begin{pmatrix} \frac{9}{2} & -\frac{9}{2} \\ -\frac{11}{2} & \frac{11}{2} \\ -\frac{13}{2} & \frac{13}{2} \\ \frac{15}{2} & -\frac{15}{2} \end{pmatrix} \begin{pmatrix} \frac{17}{2} & -\frac{17}{2} \\ -\frac{19}{2} & \frac{19}{2} \\ -\frac{21}{2} & \frac{21}{2} \\ \frac{23}{2} & -\frac{23}{2} \end{pmatrix} \begin{pmatrix} \frac{25}{2} & -\frac{25}{2} \\ -\frac{27}{2} & \frac{27}{2} \\ -\frac{29}{2} & \frac{29}{2} \\ \frac{31}{2} & -\frac{31}{2} \end{pmatrix} \begin{pmatrix} \frac{33}{2} & -\frac{33}{2} \\ -\frac{35}{2} & \frac{35}{2} \\ -\frac{37}{2} & \frac{37}{2} \\ \frac{39}{2} & -\frac{39}{2} \end{pmatrix}$$

By adding  $41/2$  to each entry of  $S$ , we obtain a 5 -tuple magic  $(4, 2)$ -rectangle  $T$  as below.

$$T := \begin{pmatrix} 21 & 20 \\ 19 & 22 \\ 18 & 23 \\ 24 & 17 \end{pmatrix} \begin{pmatrix} 25 & 16 \\ 15 & 26 \\ 14 & 27 \\ 28 & 13 \end{pmatrix} \begin{pmatrix} 29 & 12 \\ 11 & 30 \\ 10 & 31 \\ 32 & 9 \end{pmatrix} \begin{pmatrix} 33 & 8 \\ 7 & 34 \\ 6 & 35 \\ 36 & 5 \end{pmatrix} \begin{pmatrix} 37 & 4 \\ 3 & 38 \\ 2 & 39 \\ 40 & 1 \end{pmatrix}$$

Clearly, the sum over each row of any array is 41 and the sum over each column is 82.

**Proposition 4** If a balanced centrally  $n$  -tuple symmetric  $(p_1, q)$  -rectangle  $R$  and a centrally  $n$  -tuple symmetric  $(p_2, q)$  -rectangle  $S$  exist, then a centrally  $n$  -tuple symmetric  $(p_1 + p_2, q)$  -rectangle  $T$  exists. If  $S$  is a balanced rectangle, then  $T$  can also be chosen to be balanced.

**Proof.** Suppose  $S$  has type  $x$ . By Lemma 1, we know that there exists a balanced centrally  $n$  -tuple symmetric  $(p_1, q)$  -rectangle  $R'$  of type  $x + np_2q/2$ . Then by stacking  $R'$  and  $S$  together, we obtain a rectangle  $T$  whose rows' and columns' sum is zero. Thus,  $T$  is a centrally  $n$  -tuple symmetric  $(p_1 + p_2, q)$  -rectangle of type  $x$ . If  $S$  is balanced, then it is easy to see that  $T$  is also balanced.  $\square$

Since  $n$  -tuple magic  $(p, q)$  -rectangles correspond to centrally  $n$  -tuple symmetric  $(p, q)$  -rectangles of type  $-1/2$ , we have the following corollary.

**Corollary 1** Suppose an  $n$  -tuple magic  $(p_1, q)$  -rectangle and a balanced centrally  $n$  -tuple symmetric  $(p_2, q)$  -rectangle exist. Then an  $n$  -tuple magic  $(p_1 + p_2, q)$  -rectangle exists.

Using the concept of a centrally  $n$  -tuple symmetric rectangle, we can prove the existence of even  $n$  -tuple magic rectangles. Our tools are the balanced centrally  $n$  -tuple symmetric  $(2, 4)$  -rectangle  $A := (a_{i,j}^1)(a_{i,j}^2)...(a_{i,j}^n)$  given by

$$(a_{i,j}^t) = \begin{pmatrix} 4t-3 & -4t+2 & -4t+1 & 4t \\ -4t+3 & 4t-2 & 4t-1 & -4t \end{pmatrix},$$

and the  $n$  -tuple magic  $(2, 6)$  -rectangle  $B := (b_{i,j}^1)(b_{i,j}^2)...(b_{i,j}^n)$  given by

$$b_{1,j}^t = \begin{cases} 1+12(n-t) & \text{if } j=1, \\ 11+12(n-t) & \text{if } j=2, \\ 3+12(n-t) & \text{if } j=3, \\ 9+12(t-1) & \text{if } j=4, \\ 8+12(t-1) & \text{if } j=5, \\ 7+12(t-1) & \text{if } j=6, \end{cases}$$

and

$$b_{2,j}^t = \begin{cases} 12+12(t-1) & \text{if } j=1, \\ 2+12(t-1) & \text{if } j=2, \\ 10+12(t-1) & \text{if } j=3, \\ 4+12(n-t) & \text{if } j=4, \\ 5+12(n-t) & \text{if } j=5, \\ 6+12(n-t) & \text{if } j=6, \end{cases}$$

for all  $t \in \{1, 2, \dots, n\}$ .

**Proposition 5** Let  $q > 2$  be an even integer. Then an  $n$  -tuple magic  $(2, q)$  -rectangle exists.

**Proof.** We induct on  $q$ . The existence of  $n$  -tuple rectangles  $A$  and  $B$  shows that we need only prove the proposition for  $q \geq 8$ . Assume we know that an  $n$  -tuple magic  $(2, q')$  -rectangle exists for all even  $q' < q$ . Then we know an  $n$  -tuple magic  $(2, q-4)$  -rectangle  $R$  exists. By Corollary 1, we can add  $R$  and  $A$  together to form an  $n$  -tuple magic  $(2, q)$  -rectangle.  $\square$

**Proposition 6** Let  $p$  and  $q$  be even positive integers with  $(p, q) \neq (2, 2)$ . Then an  $n$  -tuple magic  $(p, q)$  -rectangle exists.

**Proof.** By Proposition 5, we can assume that  $q > 2$ . Using  $A$  and Proposition 4, induction shows that a balanced centrally  $n$  -tuple symmetric  $(p, 4)$  -rectangle  $R$  exists. Thus, an  $n$  -tuple magic  $(p, 4)$  -rectangle exists and we can assume that  $q > 4$ . Now assume that an  $n$  -tuple magic  $(p, q')$  -rectangle exists for all even  $q' < q$ . We then know that an  $n$  -tuple magic  $(p, q-4)$  -rectangle  $S$  exists. By Corollary 1, we can add  $R$  and  $S$  together to give an  $n$  -tuple magic  $(p, q)$  -rectangle.  $\square$

**Example 2** The following arrays are examples of even  $n$  -tuple magic rectangles.

A triple magic  $(6, 4)$  -rectangle

$$\begin{pmatrix} 1 & 19 & 66 & 60 \\ 50 & 68 & 17 & 11 \\ 3 & 21 & 64 & 58 \\ 70 & 52 & 9 & 15 \\ 23 & 5 & 56 & 62 \\ 72 & 54 & 7 & 13 \end{pmatrix} \begin{pmatrix} 25 & 43 & 42 & 36 \\ 26 & 44 & 41 & 35 \\ 27 & 45 & 40 & 34 \\ 46 & 28 & 33 & 39 \\ 47 & 29 & 32 & 38 \\ 48 & 30 & 31 & 37 \end{pmatrix} \begin{pmatrix} 49 & 67 & 18 & 12 \\ 2 & 20 & 65 & 59 \\ 51 & 69 & 16 & 10 \\ 22 & 4 & 57 & 63 \\ 71 & 53 & 8 & 14 \\ 24 & 6 & 55 & 61 \end{pmatrix}$$

Then each row sum of any array is 146 and each column sum of any array is 219.

A 4 -tuple magic (4, 4)-rectangle

$$\begin{pmatrix} 7 & 12 & 49 & 62 \\ 50 & 61 & 8 & 11 \\ 16 & 3 & 58 & 53 \\ 57 & 54 & 15 & 4 \end{pmatrix} \begin{pmatrix} 23 & 28 & 33 & 46 \\ 34 & 45 & 24 & 27 \\ 32 & 19 & 42 & 37 \\ 41 & 38 & 31 & 20 \end{pmatrix} \begin{pmatrix} 39 & 44 & 17 & 30 \\ 18 & 29 & 40 & 43 \\ 48 & 35 & 26 & 21 \\ 25 & 22 & 47 & 36 \end{pmatrix} \begin{pmatrix} 55 & 60 & 1 & 14 \\ 2 & 13 & 56 & 59 \\ 64 & 51 & 10 & 5 \\ 9 & 6 & 63 & 52 \end{pmatrix}$$

Then each row sum and each column sum of any array in a rectangle equals 130.

### III. The $n$ -Fold Self-Union of Complete Bipartite Graphs

For any integer  $n \geq 1$ , the  $n$ -fold self-union of a graph  $G$ , denoted by  $nG$ , is the union of  $n$  disjoint copies of  $G$ . In this section we identify the sufficient condition for degree-magic labellings of the  $n$ -fold self-union of complete bipartite graphs  $nK_{p,q} = K_{p,q}^1 \cup K_{p,q}^2 \cup \dots \cup K_{p,q}^n$  to exist.

**Theorem 3** For any integer  $n \geq 1$  and even integers  $p, q > 1$ , let  $K_{p,q}^t$  be the  $t^{\text{th}}$  copy of  $K_{p,q}$  for all  $t \in \{1, 2, \dots, n\}$ . A mapping  $f$  from  $E(nK_{p,q})$  into positive integers given by

$$f(u_i^t v_j^t) = r_{i,j}^t \text{ for every } u_i^t v_j^t \in E(K_{p,q}^t),$$

is a  $d$ -magic labelling of  $nK_{p,q}$  if and only if  $R := (r_{i,j}^1)(r_{i,j}^2) \dots (r_{i,j}^n)$  is an  $n$ -tuple magic  $(p, q)$ -rectangle.

**Proof.** Let  $U^t = \{u_1^t, u_2^t, \dots, u_p^t\}$  and  $V^t = \{v_1^t, v_2^t, \dots, v_q^t\}$  be partite sets of  $K_{p,q}^t$ . Suppose that  $R$  is an  $n$ -tuple magic  $(p, q)$ -rectangle. Then  $f$  is a bijection from  $E(nK_{p,q})$  onto  $\{1, 2, \dots, npq\}$ . For any  $u_i^t \in U^t$ , we have

$$\begin{aligned} f^*(u_i^t) &= \sum_{j=1}^q f(u_i^t v_j^t) = \sum_{j=1}^q r_{i,j}^t \\ &= \frac{q(npq+1)}{2} = \frac{npq+1}{2} \deg(u_i^t), \end{aligned}$$

and for any  $v_j^t \in V^t$ , we have

$$\begin{aligned} f^*(v_j^t) &= \sum_{i=1}^p f(u_i^t v_j^t) = \sum_{i=1}^p r_{i,j}^t \\ &= \frac{p(npq+1)}{2} = \frac{npq+1}{2} \deg(v_j^t). \end{aligned}$$

i.e.,  $f$  is a  $d$ -magic labelling of  $nK_{p,q}$ .

Now suppose that  $f$  is a  $d$ -magic labelling of  $nK_{p,q}$ . For all  $1 \leq i \neq s \leq p$ , we have

$$\begin{aligned} \sum_{j=1}^q r_{i,j}^t &= \sum_{j=1}^q f(u_i^t v_j^t) = f^*(u_i^t) \\ &= f^*(u_s^t) = \sum_{j=1}^q f(u_s^t v_j^t) = \sum_{j=1}^q r_{s,j}^t. \end{aligned} \tag{5}$$

For all  $1 \leq j \neq z \leq q$ , we have

$$\begin{aligned} \sum_{i=1}^p r_{i,j}^t &= \sum_{i=1}^p f(u_i^t v_j^t) = f^*(v_j^t) \\ &= f^*(v_z^t) = \sum_{i=1}^p f(u_i^t v_z^t) = \sum_{i=1}^p r_{i,z}^t. \end{aligned} \tag{6}$$

By (5), we have

$$\sum_{j=1}^q r_{i,j}^t = \sum_{j=1}^q r_{s,j}^t = \frac{q(npq+1)}{2}.$$

By (6), we have

$$\sum_{i=1}^p r_{i,j}^t = \sum_{i=1}^p r_{i,z}^t = \frac{p(npq+1)}{2}.$$

Therefore,  $R$  is an  $n$ -tuple magic  $(p, q)$ -rectangle.  $\square$

According to Theorem 3 and Proposition 6, we obtain the following result.

**Proposition 7** Let  $p$  and  $q$  be even positive integers with  $(p, q) \neq (2, 2)$ . Then  $nK_{p,q}$  is a  $d$ -magic graph for all integers  $n \geq 1$ .

**Example 3** We can construct a  $d$ -magic graph  $3K_{4,8}$  (see Figure 1) with the labels on edges  $u_i^t v_j^t$  of  $3K_{4,8}$ , where  $1 \leq i \leq 4$ ,  $1 \leq j \leq 8$  and  $1 \leq t \leq 3$ , in TABLE I.




Figure 1. A  $d$ -magic graph  $3K_{4,8}$ .

TABLE I. THE LABELS ON EDGES OF D-MAGIC GRAPH  $3K_{4,8}$

Vertices	$v_1^1$	$v_2^1$	$v_3^1$	$v_4^1$	$v_5^1$	$v_6^1$	$v_7^1$	$v_8^1$
$u_1^1$	1	8	93	92	9	16	85	84
$u_2^1$	6	3	90	95	14	11	82	87
$u_3^1$	91	94	4	5	83	86	12	13
$u_4^1$	96	89	7	2	88	81	15	10

Vertices	$v_1^2$	$v_2^2$	$v_3^2$	$v_4^2$	$v_5^2$	$v_6^2$	$v_7^2$	$v_8^2$
$u_1^2$	33	40	61	60	41	48	53	52
$u_2^2$	38	35	58	63	46	43	50	55
$u_3^2$	59	62	36	37	51	54	44	45
$u_4^2$	64	57	39	34	56	49	47	42

Vertices	$v_1^3$	$v_2^3$	$v_3^3$	$v_4^3$	$v_5^3$	$v_6^3$	$v_7^3$	$v_8^3$
$u_1^3$	65	72	29	28	73	80	21	20
$u_2^3$	70	67	26	31	78	75	18	23
$u_3^3$	27	30	68	69	19	22	76	77
$u_4^3$	32	25	71	66	24	17	79	74

### Acknowledgments

This work was supported by Rajamangala University of Technology Lanna and Department of Mathematics, Faculty of Science, King Mongkut's University of Technology Thonburi, Thailand.

### References

- [1] T. Harmuth, "Über magische Quadrate und ähnliche Zahlenfiguren," Arch. Math. Phys., vol. 66, pp. 286-313, 1881.
- [2] T. Harmuth, "Über magische Rechtecke mit ungeraden Seitenzahlen," Arch. Math. Phys., vol. 66, pp. 413-447, 1881.
- [3] R. Sun, "Existence of magic rectangles," Nei Mongol Daxue Xuebao Ziran Kexue, vol. 21, no. 1, pp. 10-16, 1990.
- [4] T. Bier and G. Rogers, "Balanced magic rectangles," European J. Combin., vol. 14, pp. 285-299, 1993.
- [5] T. Bier and A. Kleinschmidt, "Centrally symmetric and magic rectangles," Discrete Math., vol. 176, pp. 29-42, 1997.
- [6] T. Hagedorn, "Magic rectangles revisited," Discrete Math., vol. 207, pp. 65-72, 1999.
- [7] T. Hagedorn, "On the existence of magic  $n$ -dimensional rectangles," Discrete Math., vol. 207, pp. 53-63, 1999.
- [8] J. Sedláček, "Theory of graphs and its applications," Proc. Symp. Smolenice, Problem 27, Praha, pp. 163-164, 1963.
- [9] B.M. Stewart, "Magic graphs," Canad. J. Math., Vol.18, pp. 1031-1059, 1966.

- [10] J.A. Gallian, "A dynamic survey of graph labeling," Electron. J. Combin., #DS6, vol.16, 2009.
- [11] L'. Bezegová and J. Ivančo, "An extension of regular supermagic graphs," Discrete Math., vol. 310, pp. 3571-3578, 2010.