

Analytical applications on the Dunkl-type Fock space $F_k(C^d)$

Fethi Soltani and Akram Nemri

Abstract— In this work, we establish Heisenberg-type uncertainty principle for the Dunkl-type Fock space $F_k(C^d)$. Next, we give an application of the classical theory of reproducing kernels to the Tikhonov regularization problem for operator $L: F_k(C^d) \rightarrow H$, where H be a Hilbert space. Finally, we come up with some results regarding the Tikhonov regularization problem and the Heisenberg-type uncertainty principle for the Dunkl-type Segal-Bargmann transform B_k .

Keywords—component, Dunkl-type Fock spaces, extremal functions, Tikhonov regularization, Heisenberg-type uncertainty principle.

I- Introduction

In this work, by using the theory of operators on Hilbert spaces [3, 4] we establish Heisenberg-type uncertainty principle for the Dunkl-type Fock space $F_k(C^d)$ [1, 11]. In the Dunkl setting, the Heisenberg-type uncertainty principle for the Dunkl transform D_k [2,5], is studied by Rosler in [7] and by Soltani in [15]. This is an important support for show in this work, the Heisenberg-type uncertainty principle for the Dunkl-type Segal-Bargmann transform B_k [1, 11]. Next, building on the ideas of Matsuura et al. [6], Saitoh [10], and Soltani [12], we give an application of the classical theory of reproducing kernels [9] to the Tikhonov regularization problem on the Dunkl-type Fock space $F_k(C^d)$. More precisely, let $L: F_k(C^d) \rightarrow H$ be a bounded operator from $F_k(C^d)$ into a Hilbert space H ,

for all $\lambda > 0$, $h \in H$, the infimum $\inf_{f \in F_k(C^d)} (\lambda \|f\|_{F_k(C^d)}^2 + \|h - Lf\|_H^2)$, is attained at one function $f_{\lambda,h}^*$, called the extremal function. As applications, we study the extremal functions associated to the following operators:

- the operator $Lf(w) = (B_k)^{-1} f(w)$,
- the operator $Lf(w) = (D_k)^{-1} (B_k)^{-1} f(w)$,
- and the operator $Lf(w) = f(-iw)$.

In the Dunkl setting, the extremal functions are studied by Soltani in several directions, where L is the Dunkl-type Weierstrass transform [12], L is the Dunkl multiplier operators [13,16] and L is the dual Dunkl-Sonine operator [14].

The contents of the paper are as follows. In Section 2, we recall some properties of the Dunkl-type Fock space $F_k(C^d)$ and we establish Heisenberg-type uncertainty principle for this space. In Section 3, we give an application of the classical theory of reproducing kernels to the Tikhonov regularization problem for bounded operator $L: F_k(C^d) \rightarrow H$, where H be a Hilbert space. The last section is devoted to give some results concerning the Tikhonov regularization problem and the Heisenberg-type uncertainty principle for the Dunkl-type Segal-Bargmann transform B_k .

II- Dunkl-type Segal-Bargmann space

The Dunkl operators $T_j(k)$; $j=1, \dots, d$, on R^d associated with the positive root system \mathfrak{R}^+ and the multiplicity function k are given, for a function f of class C^1 on R^d , by

$$T_j(k)f(x) := \frac{\partial}{\partial x_j} f(x) + \sum_{\alpha \in \mathfrak{R}^+} k(\alpha) \alpha_j \frac{f(x) - f(\sigma_\alpha x)}{\langle \alpha, x \rangle}$$

where $\sigma_\alpha x := x - \frac{2\langle \alpha, x \rangle}{|\alpha|^2} \alpha$.

Fethi Soltani
 Department of Mathematics
 Faculty of Science, Jazan University
 Jazan, Saudi Arabia

Akram Nemri
 Department of Mathematics
 Faculty of Science, Jazan University
 Jazan, Saudi Arabia

For $y \in \mathbb{R}^d$, the initial problem $T_j(k)u(.,y)(x) = y_j u(x,y)$, $j=1, \dots, d$, with $u(0,y) = 1$ admits a unique analytic solution on \mathbb{R}^d , which will be denoted by $E_k(x,y)$ and called Dunkl kernel [2]. This kernel has a unique analytic extension to $\mathbb{C}^d \times \mathbb{C}^d$ (see [8]).

For $z, w \in \mathbb{C}^d$, we define $K(z, w) := E_k(z, \bar{w})$. From the properties of the Dunkl kernel [2, 8], the kernel K is continuous and the function $z \rightarrow K(z, .)$ is holomorphic for all $w \in \mathbb{C}^d$. Further, $K(z, w) = \overline{K(w, z)}$ and $K(z, w)$ is a positive definite kernel. Then, there exists a Hilbert space $F_k(\mathbb{C}^d)$ of holomorphic functions on \mathbb{C}^d such that K is its reproducing kernel. The Hilbert space $F_k(\mathbb{C}^d)$ contains the \mathbb{C} -algebra $P(\mathbb{C}^d)$ of polynomial functions on \mathbb{C}^d as a dense subspace. In particular, if we denote by $\langle \cdot, \cdot \rangle_{F_k(\mathbb{C}^d)}$ the inner product in $F_k(\mathbb{C}^d)$, then

$$\langle p, q \rangle_{F_k(\mathbb{C}^d)} = p(T(k)) \overline{q(\bar{z})}_{z=0}, p, q \in P(\mathbb{C}^d),$$

where $p(T(k))$ is the operator formed by replacing z_j by $T_j(k)$ for $j=1, \dots, d$. We shall call $F_k(\mathbb{C}^d)$ the Dunkl-type Fock space.

III- Heisenberg-type uncertainty principle

We consider the following general result from functional analysis.

Theorem 3.1. (See [3, 4]). Let A and B be self adjoint operators on a Hilbert space H . We define the commutator of A and B to $[A, B] := AB - BA$. Then we have that

$$\|(A-a)f\|_H \|(B-b)f\|_H \geq \frac{1}{2} |\langle [A, B]f, f \rangle_H|, \text{ for all } f \in \text{Dom}(AB) \cap \text{Dom}(BA), \text{ and all } a, b \in \mathbb{R}.$$

By applying Theorem 3.1 we establish the Heisenberg-type uncertainty principle for the Dunkl-type Fock space $F_k(\mathbb{C}^d)$.

Theorem 3.2. Let $f \in F_k(\mathbb{C}^d)$. For all $a, b \in \mathbb{R}$, we have

$$\|(T_j(k) + z_j - a)f\|_{F_k(\mathbb{C}^d)} \|(T_j(k) - z_j + ib)f\|_{F_k(\mathbb{C}^d)} \geq N_k(f)$$

$$\text{where } N_k(f) = \left\| \|f\|_{F_k(\mathbb{C}^d)}^2 + \langle P_j(k)f, f \rangle_{F_k(\mathbb{C}^d)} \right\|, \text{ and } P_j(k)f(z) = \sum_{\alpha \in \mathbb{N}^+} k(\alpha) (\alpha_j)^2 f(\sigma_\alpha z).$$

This uncertainty principle can be written as the following.

Theorem 3.3. Let $f \in F_k(\mathbb{C}^d)$. Then

$$M_k^+(f) M_k^-(f) \geq \|f\|_{F_k(\mathbb{C}^d)}^2 (N_k(f))^2$$

where

$$M_k^\pm(f) = \|f\|_{F_k(\mathbb{C}^d)}^2 \|A_k^\pm(f)\|_{F_k(\mathbb{C}^d)}^2 - \left| \langle A_k^\pm(f), f \rangle_{F_k(\mathbb{C}^d)} \right|^2$$

$$A_k^\pm(f) = (T_j(k) \pm z_j) f.$$

IV- Extremal functions on $F_k(\mathbb{C}^d)$

Let $\lambda > 0$ and let $L: F_k(\mathbb{C}^d) \rightarrow H$ be a bounded linear operator from $F_k(\mathbb{C}^d)$ into a Hilbert space H . We denote by $\langle f, g \rangle_{L, \lambda}$ the inner product defined on the space $F_k(\mathbb{C}^d)$ by

$$\langle f, g \rangle_{L, \lambda} := \lambda \langle f, g \rangle_{F_k(\mathbb{C}^d)} + \langle Lf, Lg \rangle_H.$$

By using the theory of extremal function and reproducing kernel of Hilbert space [9,10] we establish the extremal function associated to the operator L on the Dunkl-type Fock space $F_k(\mathbb{C}^d)$.

Theorem 4.1. Let $\lambda > 0$. The Fock space $(F_k(\mathbb{C}^d), \langle \cdot, \cdot \rangle_{L, \lambda})$ possesses a reproducing kernel $K_{L, \lambda}(w, z); w, z \in \mathbb{C}^d$, which satisfying the equation $(\lambda I + L^*L)K_{L, \lambda}(w, .) = K(w, .)$. Moreover the kernel $K_{L, \lambda}(w, z)$ satisfies the following properties

- (i) $\|K_{L, \lambda}(w, .)\|_{F_k(\mathbb{C}^d)} \leq \frac{\sqrt{E_k(w, \bar{w})}}{\lambda}$,
- (ii) $\|LK_{L, \lambda}(w, .)\|_H \leq \sqrt{\frac{E_k(w, \bar{w})}{\lambda}}$,
- (iii) $\|L^*LK_{L, \lambda}(w, .)\|_{F_k(\mathbb{C}^d)} \leq \sqrt{E_k(w, \bar{w})}$.

From this theorem we obtain the main result of this section.

Theorem 4.2. For any $h \in H$ and for any $\lambda > 0$, there exists a unique function $f_{\lambda, h}^*$, where the infimum

$$\inf_{f \in F_k(\mathbb{C}^d)} \left(\lambda \|f\|_{F_k(\mathbb{C}^d)}^2 + \|h - Lf\|_H^2 \right) \text{ is attained.}$$

Moreover, the extremal function $f_{\lambda, h}^*$ is given by

$f_{\lambda,h}^*(w) = \langle h, LK_{L,\lambda}(w, \cdot) \rangle_H$ and satisfies the inequality

$$|f_{\lambda,h}^*(w)| \leq \sqrt{\frac{E_k(w, w)}{\lambda}} \|h\|_H.$$

v. Dunkl-type Segal-Bargmann transform

Let ω_k denote the weight function

$$\omega_k(y) := \prod_{\alpha \in \mathfrak{R}^+} |\langle \alpha, y \rangle|^{2k(\alpha)}, \quad y \in \mathbb{R}^d.$$

Let c_k be the Mehta-type constant given by

$$c_k := \left(\int_{\mathbb{R}^d} e^{-|y|^2} \omega_k(y) dy \right)^{-1}.$$

We denote by μ_k the measure on \mathbb{R}^d given by

$d\mu_k(y) := c_k \omega_k(y) dy$, and by $L_k^2(\mathbb{R}^d)$, the space of measurable functions f on \mathbb{R}^d , such that

$$\|f\|_{L_k^2(\mathbb{R}^d)} := \left(\int_{\mathbb{R}^d} |f(y)|^2 dy \right)^{\frac{1}{2}} < \infty.$$

We denote by U_k the kernel given for $z, w \in \mathbb{C}^d$, by

$$U_k(z, w) := e^{-\frac{\langle z, z \rangle + \langle w, w \rangle}{2}} E_k(\sqrt{2}z, w).$$

The kernel U_k gives rise to an integral transform

B_k , which is called Dunkl-type Segal-Bargmann transform on

\mathbb{C}^d , and defined for f in $L_k^2(\mathbb{R}^d)$, by

$$B_k(f)(z) := \int_{\mathbb{R}^d} U_k(z, x) f(x) d\mu_k(x), \quad z \in \mathbb{C}^d.$$

The transform B_k is an isometric isomorphism of

$L_k^2(\mathbb{R}^d)$ onto $F_k(\mathbb{C}^d)$, and for $f \in L_k^2(\mathbb{R}^d)$ we have

$$\|B_k(f)\|_{F_k(\mathbb{C}^d)} = \|f\|_{L_k^2(\mathbb{R}^d)},$$

$$B_k D_k(f)(w) = B_k(f)(-iw), \quad w \in \mathbb{C}^d,$$

where D_k is the Dunkl transform [2,5].

Application 5.1. (i) If $H = L_k^2(\mathbb{R}^d)$ and $L = (B_k)^{-1}$,

then $L^* = B_k$ and $L^*L = I$. Thus, by Theorem 4.1, we

deduce that $K_{L,\lambda}(w, z) = \frac{1}{\lambda+1} K(w, z)$, and for

$$h \in L_k^2(\mathbb{R}^d), \quad f_{\lambda,h}^*(w) = \frac{1}{\lambda+1} B_k(h)(w).$$

(ii) If $H = F_k(\mathbb{C}^d)$ and $Lf(w) = f(-iw)$, then

$L^*f(w) = f(iw)$ and $L^*L = I$. Thus, by Theorem 4.1, we

deduce that $K_{L,\lambda}(w, z) = \frac{1}{\lambda+1} K(w, z)$, and for

$$h \in F_k(\mathbb{C}^d), \quad f_{\lambda,h}^*(w) = \frac{1}{\lambda+1} h(iw).$$

Application 5.2. By applying Heisenberg-type uncertainty principle for the Dunkl transform ([15], Theorem 2), we deduce the following Heisenberg-type uncertainty principle for the Dunkl-type Segal-Bargmann transform B_k .

Let $a > 0$. There exists $C_a > 0$ such that for all

$f \in L_k^2(\mathbb{R}^d)$ we have

$$\left\| e^{-\frac{|x|^2}{2}} f \right\|_{L_k^2(\mathbb{R}^d)} \leq C_a \left\| |x|^a e^{-\frac{|x|^2}{2}} f \right\|_{L_k^2(\mathbb{R}^d)} \left[\sum_{j=1}^d \|z_j B_k(f)\|_{F_k(\mathbb{C}^d)}^2 \right]^{\frac{1}{2a+2}}$$

Acknowledgment

This work was supported by the Deanship of the Scientific Research of Jazan University, Saudi Arabia (Project N^o 588-Sabic 2 - 36).

References

- [1] S. Ben Said and B. Ørsted, Segal-Bargmann transforms associated with Coxeter groups, *Math. Ann.* **334** (2006) 281-323.
- [2] C.F. Dunkl, Hankel transforms associated to finite reflection groups, *Contemp. Math.* **138** (1992) 123-138.
- [3] G. Folland, *Harmonic analysis on phase space*, Annals of Mathematics Studies 122, Princeton University Press, Princeton, New Jersey, 1989.
- [4] K. Gröchenig, *Foundations of time-frequency analysis*, Birkhauser, Boston, 2001.
- [5] M.F.E.de Jeu, The Dunkl transform, *Inv. Math.* **113** (1993) 147-162.
- [6] T. Matsuura, S. Saitoh and D.D. Trong, Inversion formulas in heat conduction multidimensional spaces, *J. Inv. Ill-posed Problems* **13** (2005) 479-493.
- [7] M. Rösler, An uncertainty principle for the Dunkl transform, *Bull. Austral. Math. Soc.* **59** (1999) 353-360.
- [8] E.M. Opdam, Dunkl operators, Bessel functions and the discriminant of a finite Coxeter group, *Compositio Math.* **85** (3) (1993) 333-373.
- [9] S. Saitoh, Hilbert spaces induced by Hilbert space valued functions, *Proc. Amer. Math. Soc.* **89** (1983) 74-78.
- [10] S. Saitoh, Best approximation, Tikhonov regularization and reproducing kernels, *Kodai Math. J.* **28** (2005) 359-367.
- [11] F. Soltani, Generalized Fock spaces and Weyl commutation relations for the Dunkl kernel, *Pacific J. Math.* **214** (2004) 379-397.

- [12] F. Soltani, Inversion formulas in the Dunkl-type heat conduction on \mathbb{R}^d , Appl. Anal. **84** (2005) 541-553.
- [13] F. Soltani, Best approximation formulas for the Dunkl L^2 -multiplier operators on \mathbb{R}^d , Rocky Mountain J. Math. **42** (2012) 305-328.
- [14] F. Soltani, Multiplier operators and extremal functions related to the dual Dunkl-Sonine operator, Acta Math. Sci. **33B**(2) (2013) 430-442.
- [15] F. Soltani, A general form of Heisenberg-Pauli-Weyl uncertainty inequality for the Dunkl transform, Int. Trans. Spec. Funct. **24** (5) (2013) 401-409.
- [16] F. Soltani, Dunkl multiplier operators and applications, Int. Trans. Spec. Funct. **25** (11) (2014) 898-908.